


**LECTURE NOTES ON ENGINEERING MATHEMATIC-I**

**(COMMON FOR ALL BRANCH)**



**PREPARED BY,**

**SAMIRA KUMAR PATHI  
LECTURER IN MATHEMATICS,  
GOVT. POLYTECHNIC, GAJAPATI**

**MATHEMATICS –I FOR DIPLOMA 1<sup>ST</sup> SEM STUDENTS**

(Prepared by Sri Samira Kumar Pathi ,Lecturer in Mathematics, G.P.Gajapati)

## MATRIX

A *matrix* is a rectangular array of numbers (or other mathematical objects) arranged in rows and columns for which operations such as addition and multiplication are defined.

$$\begin{bmatrix} a & d \\ b & e \\ c & f \end{bmatrix}; \begin{bmatrix} 1 & 2 & 3 \\ 5 & 7 & 8 \end{bmatrix}; \begin{bmatrix} 1 & 2 & 5 \\ 8 & 9 & 10 \\ 7 & 5 & 15 \end{bmatrix}; \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

### Uses of matrix in real life:-

This is a football match. Here A passes the ball to B, B passes the ball to A, B passes the ball to C and C passes the ball to D. This can inform to the computer using matrix as follows

	A	B	C	D
A	0	1	0	0
B	1	0	1	0
C	0	0	0	1
D	0	0	0	0

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

### Order of a matrix:-

The number of rows and columns that a matrix has is called its order. By convention, rows are listed first; and columns, second. Thus, we would say that the order (or dimension) of the matrix below is 3 x 4, meaning that it has 3 rows and 4 columns.



$$\begin{bmatrix} 1 & 2 & 2 & 5 \\ 8 & 9 & 1 & 10 \\ 7 & 5 & 9 & 15 \end{bmatrix}$$

The matrix with m rows and n column is m x n (m-by-n)

Types of matrix:-

Square matrix:-

If a matrix have same number of rows and columns.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \quad \begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & 3 \\ 2 & 0 & 4 \end{bmatrix}$$

Null matrix/zero matrix:-

If all the elements of a matrix are zero then the matrix is called a zero/ null matrix.

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \quad \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Row matrix & Column Matrix:-

If a matrix has only one row (or column) is called a row matrix (or column matrix)

$$\begin{bmatrix} 1 & 1 & 2 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

Diagonal Matrix:-

If all the non-diagonal entries of a square matrix are zero and at least one diagonal entry is non-zero then the matrix is called a Diagonal matrix.

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 6 \end{bmatrix} \quad \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

**Properties of determinant:-**

- i) If any two rows or columns of a matrix A are equal then the determinant of A is zero.

Ex:-  $\begin{vmatrix} 2 & 2 & 0 \\ 1 & 1 & 1 \\ 1 & 1 & 5 \end{vmatrix}$  (calculate the determinant) =0

$$\begin{vmatrix} 2 & 1 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 5 \end{vmatrix} \text{ (Calculate the determinant)=0}$$

$$\begin{vmatrix} 2 & 1 & 1 \\ 1 & 1 & 1 \\ 2 & 1 & 1 \end{vmatrix} \text{ (Calculate the determinant)=0}$$

ii) If we exchange any two rows or columns then the absolute value of the determinant does not change. But the sign of the determinant is changed.

Ex:-  $\begin{vmatrix} 5 & 1 & 2 \\ 1 & 1 & 1 \\ 2 & 1 & 1 \end{vmatrix}$  (Calculate the determinant) = -1

Exchanging 1<sup>st</sup> row and 2<sup>nd</sup> row

$|A| = \begin{vmatrix} 1 & 1 & 1 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix}$  (Calculate the determinant) = +1

iii) if  $k$  is multiplied to each element of any row or column of a matrix  $A$  then the determinant of the matrix so formed is  $k \cdot \det(A)$

Ex-  $|B| = \begin{vmatrix} 2 & 2 & 2 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix}$  (Calculate the determinant) = +2

$$\begin{vmatrix} 2 & 2 & 2 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix} = 2(1 \cdot 1 - 1 \cdot 2) - 2(5 \cdot 1 - 2 \cdot 2) + 2(5 \cdot 1 - 2 \cdot 1) = -2 - 2 + 6 = 2 = 2 \cdot 1 = 2 \cdot \begin{vmatrix} 1 & 1 & 1 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix}$$

$|B| = 2 \cdot |A|$

Ex-  $\begin{vmatrix} 2 & 2 & 2 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix} = \begin{vmatrix} 1+1 & 1+1 & 1+1 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 1 & 1 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 1 & 1 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix}$

$$\begin{vmatrix} 5 & 2 & 3 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix} = \begin{vmatrix} 2+3 & 1+1 & 2+1 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix} = \begin{vmatrix} 2 & 1 & 2 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix} + \begin{vmatrix} 3 & 1 & 1 \\ 5 & 1 & 2 \\ 2 & 1 & 1 \end{vmatrix}$$

$$\begin{vmatrix} a+b & c+d & e+f \\ g & h & i \\ j & k & l \end{vmatrix} = \begin{vmatrix} a & c & e \\ g & h & i \\ j & k & l \end{vmatrix} + \begin{vmatrix} b & d & f \\ g & h & i \\ j & k & l \end{vmatrix}$$

### PROBLEM

I)  $\begin{vmatrix} 3 & 6 & 9 \\ 1 & 1 & 1 \\ 4 & 8 & 12 \end{vmatrix}$  (without expanding determinant and use the properties of determinant)

$$\begin{vmatrix} 3.1 & 3.2 & 3.3 \\ 1 & 1 & 1 \\ 4.1 & 4.2 & 4.3 \end{vmatrix} \quad (\text{using 3rd property})$$

$$= 3.4 \begin{vmatrix} 1 & 2 & 3 \\ 1 & 1 & 1 \\ 1 & 2 & 3 \end{vmatrix} = 12 \times 0 \quad (\text{using 1st property}) \quad (\text{as the elements of 1st row and 3rd row are equal})$$

$$\begin{vmatrix} 12 & 3 & 9 \\ 1 & 1 & 1 \\ 8 & 8 & 4 \end{vmatrix} = 3.4 \begin{vmatrix} 4 & 1 & 3 \\ 1 & 1 & 1 \\ 2 & 2 & 1 \end{vmatrix} = 12$$

#### PROPERTY 4

If to each element of any row or column of a determinant the equimultiples of corresponding elements of other row(or column) are added, then the value of the determinant remains the same, operation is  $R_1$  by  $R_1 + kR_2$  or  $C_1$  by  $C_1 + kC_2$

**2<sup>ND</sup> PROBLEM**  $\begin{vmatrix} y+z & z & y \\ z & z+x & x \\ y & x & x+y \end{vmatrix} = 4xyz$  (without expanding evaluate the determinant)

$$= \begin{vmatrix} (y+z) + z + y & z + (z+x) + x & y + x + (x+y) \\ z & z+x & x \\ y & x & x+y \end{vmatrix} \quad (R_1 \text{ BY } R_1 + R_2 + R_3)$$

$$= \begin{vmatrix} 2(y+z) & 2(z+x) & 2(x+y) \\ z & z+x & x \\ y & x & x+y \end{vmatrix}$$

$$2 \begin{vmatrix} (y+z) & (z+x) & (x+y) \\ z & z+x & x \\ y & x & x+y \end{vmatrix}$$

$$2 \begin{vmatrix} (y+z) - ((z+x) + (x+y)) & (z+x) & (x+y) \\ z - ((z+x) + x) & z+x & x \\ y - (x + (x+y)) & x & x+y \end{vmatrix} \quad \{C_1 \rightarrow C_1 - (C_2 + C_3)\}$$

$$2 \begin{vmatrix} (y+z) - z - x - x - y & (z+x) & (x+y) \\ z - (z+x) - x & z+x & x \\ y - x - x - y & x & x+y \end{vmatrix}$$

$$2 \begin{vmatrix} -X - X & (z + x) & (x + y) \\ (-X) - X & z + x & x \\ -X - X & x & x + y \end{vmatrix}$$

$$= 2 \begin{vmatrix} -2X & (z + x) & (x + y) \\ -2X & z + x & x \\ -2X & x & x + y \end{vmatrix}$$

$$= 2X - 2x \begin{vmatrix} 1 & (z + x) & (x + y) \\ 1 & z + x & x \\ 1 & x & x + y \end{vmatrix}$$

$$= -4x \begin{vmatrix} 1 & (z + x) & (x + y) \\ 0 & 0 & -y \\ 0 & -z & 0 \end{vmatrix} \quad (\text{R2 by R2-R1, R3 by R3-R1})$$

$$= -4x \times 1 \begin{vmatrix} 0 & -y \\ -z & 0 \end{vmatrix}$$

$$= -4x \times (0 - (zy))$$

$$= 4xyz$$

2) Prove without expanding  $\begin{vmatrix} bc & a & a^2 \\ ca & b & b^2 \\ ab & c & c^2 \end{vmatrix} = \begin{vmatrix} 1 & a^2 & a^3 \\ 1 & b^2 & b^3 \\ 1 & c^2 & c^3 \end{vmatrix}$

$$\begin{vmatrix} bc & a & a^2 \\ ca & b & b^2 \\ ab & c & c^2 \end{vmatrix} = \frac{1}{a} * \frac{1}{b} * \frac{1}{c} \begin{vmatrix} bc * a & a * a & a^2 * a \\ ca * b & b * b & b^2 * b \\ ab * c & c * c & c^2 * c \end{vmatrix} \quad (\text{multiply 1st row by a, 2nd row by b, 3rd row by c})$$

$$\text{R1} \rightarrow a\text{R1}; \text{R2} \rightarrow b\text{R2}; \text{R3} \rightarrow c\text{R3}$$

$$= \frac{1}{abc} \begin{vmatrix} abc & a^2 & a^3 \\ abc & b^2 & b^3 \\ abc & c^2 & c^3 \end{vmatrix} = \frac{1}{abc} abc \begin{vmatrix} 1 & a^2 & a^3 \\ 1 & b^2 & b^3 \\ 1 & c^2 & c^3 \end{vmatrix} = 1 \cdot \begin{vmatrix} 1 & a^2 & a^3 \\ 1 & b^2 & b^3 \\ 1 & c^2 & c^3 \end{vmatrix} = \begin{vmatrix} 1 & a^2 & a^3 \\ 1 & b^2 & b^3 \\ 1 & c^2 & c^3 \end{vmatrix}$$

### Transpose of a matrix:-

If A is a matrix then the transpose of A which is denoted by  $A^T$  is formed by exchanging elements of rows by elements of columns.

$$A = \begin{vmatrix} 1 & 2 \\ 3 & 1 \end{vmatrix} \quad A^T = \begin{vmatrix} 1 & 3 \\ 2 & 1 \end{vmatrix}$$

$$A = \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix} \quad \text{then } A^T = \begin{vmatrix} a & x & p \\ b & y & q \\ c & z & r \end{vmatrix}$$

$$\text{Det}(A) = \text{det}(A^T)$$

### Minor and Co-factor , ADJ A , INVERSE OF A :-

Suppose A be a matrix and its determinant be  $\text{det}(A)$ . Then the minor of an element is the determinant of the matrix so formed after deleting the corresponding row and column to which the element lie.

$$C_{ij} = (-1)^{i+j} M_{ij}$$

$$A = \begin{vmatrix} 1 & -2 \\ 4 & 3 \end{vmatrix} \quad M_{11} = 3 \quad C_{11} = (-1)^2 \times 3 = 3 \quad |A| = 3 - (-8) = 11$$

$$M_{12} = 4 \quad C_{12} = (-1)^3 \times 4 = -4 \quad \text{ADJ } A = \begin{vmatrix} 3 & 2 \\ -4 & 1 \end{vmatrix}$$

$$M_{21} = -2 \quad C_{21} = (-1)^3 \times -2 = 2$$

$$M_{22} = 1 \quad C_{22} = (-1)^4 \times 1 = 1$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{vmatrix} 3 & 2 \\ -4 & 1 \end{vmatrix}}{11} = \begin{vmatrix} 3/11 & 2/11 \\ -4/11 & 1/11 \end{vmatrix}$$

$$A = \begin{vmatrix} 2 & -3 & 5 \\ 6 & 0 & 4 \\ 1 & 5 & -7 \end{vmatrix} \quad |A| = 2(-20) - (-3)(-46) + 5(30) = -40 - 138 + 150 = -178 + 150 = -28$$

$$M_{11} = \begin{vmatrix} 0 & 4 \\ 5 & -7 \end{vmatrix} = 0 - 20 = -20 \quad C_{11} = (-1)^2 \times -20 = -20$$

$$M_{12} = -42 - 4 = -46 \quad C_{12} = (-1)^3 \times -46 = 46$$

$$M_{13} = 30 - 0 = 30 \quad C_{13} = (-1)^4 \times 30 = 30$$

$$M_{21} = 21 - 25 = -4 \quad C_{21} = (-1)^3 \times -4 = 4$$

$$M_{22} = -14 - 5 = -19 \quad C_{22} = (-1)^4 \times -19 = -19$$

$$M_{23} = 10 - (-3) = 13 \quad C_{23} = (-1)^5 \times 13 = -13$$

$$M_{31} = -12 - 0 = -12 \quad C_{31} = (-1)^4 \times -12 = -12$$

$$M_{32} = 8 - 30 = -22$$

$$C_{32} = (-1)^5 \times -22 = 22$$

$$M_{33} = 0 - (-18) = 18$$

$$C_{33} = (-1)^6 \times 18 = 18$$

$$\text{Adj } A = \begin{bmatrix} C_{11} & C_{21} & C_{31} \\ C_{12} & C_{22} & C_{32} \\ C_{13} & C_{23} & C_{33} \end{bmatrix} = \begin{bmatrix} -20 & 4 & -12 \\ 46 & -19 & 22 \\ 30 & -13 & 18 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} -20 & 4 & -12 \\ 46 & -19 & 22 \\ 30 & -13 & 18 \end{bmatrix}}{-28}$$

### MATRIX ADDITION SUBTRACTION MULTIPLICATION

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix} \quad A + B = \begin{bmatrix} 1+1 & 2+1 \\ 3+2 & 4+1 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & 5 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix} \quad A - B = \begin{bmatrix} 1-1 & 2-1 \\ 3-2 & 4-1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 3 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix} \quad A \cdot B = \begin{bmatrix} 1 \times 1 + 2 \times 2 & 1 \times 1 + 2 \times 1 \\ 3 \times 1 + 4 \times 2 & 3 \times 1 + 4 \times 1 \end{bmatrix} = \begin{bmatrix} 5 & 3 \\ 11 & 7 \end{bmatrix}$$

### ADDITION, SUBTRACTION = ORDER SAME

### MULTIPLICATION $A = m \times n$ $B = n \times p$ $AB = M \times N$

$$A = \begin{bmatrix} 1 & 2 \\ 1 & 1 \\ 4 & 2 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 \\ 1 & 0 \\ 5 & 4 \end{bmatrix} \quad C = \begin{bmatrix} 2 & 3 & 2 \\ 1 & 1 & 0 \end{bmatrix} \quad A = 3 \times 2 \quad B = 3 \times 2 \quad C = 2 \times 3$$

$$A + B = \begin{bmatrix} 1 & 2 \\ 1 & 1 \\ 4 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 0 \\ 5 & 4 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 2 & 1 \\ 9 & 6 \end{bmatrix}$$

$$A - B = \begin{bmatrix} 1 & 2 \\ 1 & 1 \\ 4 & 2 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & 0 \\ 5 & 4 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 1 \\ -1 & -2 \end{bmatrix}$$

$$A \cdot C = \begin{bmatrix} 1 & 2 \\ 1 & 1 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} 2 & 3 & 2 \\ 1 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 \times 2 + 2 \times 1 & 1 \times 3 + 2 \times 1 & 1 \times 2 + 2 \times 0 \\ 1 \times 2 + 1 \times 1 & 1 \times 3 + 1 \times 1 & 1 \times 2 + 1 \times 0 \\ 4 \times 2 + 2 \times 1 & 4 \times 3 + 2 \times 1 & 4 \times 2 + 2 \times 0 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 5 & 2 \\ 3 & 4 & 2 \\ 10 & 14 & 8 \end{bmatrix}$$



C.A

$$= \begin{bmatrix} 2 & 3 & 2 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 1 \\ 4 & 2 \end{bmatrix} = \begin{bmatrix} 2X1 + 3X1 + 2X4 & 2X2 + 3X1 + 2X2 \\ 1X1 + 1X1 + 0X4 & 1X2 + 1X1 + 0X2 \end{bmatrix} = \begin{bmatrix} 13 & 11 \\ 2 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & x \\ 2 & -2 \end{bmatrix} + \begin{bmatrix} 3 & 4 \\ 3 & 6 \end{bmatrix} = \begin{bmatrix} 4 & 3 \\ 5 & 4 \end{bmatrix}.$$

FIND THE VALUE OF x?

$$x+4=3$$

$$x = -1 \quad \mathbf{a/b = a \cdot b^{-1}}$$

Solution of system of linear equation by Cramer's method:-

Linear equation:- A linear equation is an equation consisting of only the **variable** with power one/ degree one.

System of linear equation:- Combination of two or more linear equations is known as system of linear equation.

Suppose  $2x-y+3z=3$ ;  $x+2y+9z=2$ ;  $x-5y+3z=6$  is the system of linear equation with **variables** x,y & z each variable has degree 1.

$$2x-y+3z=3$$

$$x+2y+9z=2$$

$$x-5y+3z=6$$

$$AX=B$$

$$\equiv \begin{pmatrix} 2 & -1 & 3 \\ 1 & 2 & 9 \\ 1 & -5 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 6 \end{pmatrix}$$

$$|A| = 2(6+45)+1(3-9)+3(-5-2)=102-6-21=75$$

$$A_x = \begin{pmatrix} 3 & -1 & 3 \\ 2 & 2 & 9 \\ 6 & -5 & 3 \end{pmatrix}$$

$$|A_x| = 3(6+45)+1(6-54)+3(-10-12)=153-48-66=39$$

$$A_y = \begin{pmatrix} 2 & 3 & 3 \\ 1 & 2 & 9 \\ 1 & 6 & 3 \end{pmatrix}$$

$$|A_y| = 2(6-54) - 3(3-9) + 3(6-2) = -96 + 18 + 12 = -66$$

$$A_z = \begin{pmatrix} 2 & -1 & 3 \\ 1 & 2 & 2 \\ 1 & -5 & 6 \end{pmatrix}$$

$$|A_z| = 2(12+10) + 1(6-2) + 3(-5-2) = 44 + 4 - 21 = 48 - 21 = 27$$

$$x = \frac{|A_x|}{|A|} = 39/75$$

$$y = \frac{|A_y|}{|A|} = -66/75$$

$$z = \frac{|A_z|}{|A|} = 27/75$$

problem 2-  $2x + y = 1$  solve by creamer's rule

$$3x - y = 4$$

$$AX = B$$

$$= \begin{bmatrix} 2 & 1 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} X \\ Y \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}$$

$$|A| = -2 - 3 = -5$$

$$|A_x| = \begin{bmatrix} 1 & 1 \\ 4 & -1 \end{bmatrix} = -1 - 4 = -5$$

$$|A_y| = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} = 8 - 3 = 5$$

$$x = \frac{|A_x|}{|A|} = -5/-5 = 1$$

$$y = \frac{|A_y|}{|A|} = 5/-5 = -1$$

pr -ii  $2x + y = 1$  by matrix inversion method

$$3x - y = 4$$

$$AX = B$$

$$X = A^{-1}B$$

$$= \begin{bmatrix} 2 & 1 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} X \\ Y \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}$$

$$A = \begin{bmatrix} 2 & 1 \\ 3 & -1 \end{bmatrix} \quad |A| = -2 - 3 = -5$$

$$\begin{array}{ll} M_{11} = -1 & C_{11} = -1 \\ M_{12} = 3 & C_{12} = -3 \\ M_{21} = 1 & C_{21} = -1 \\ M_{22} = 2 & C_{22} = 2 \end{array}$$

$$\text{Adj } A = \begin{bmatrix} -1 & -1 \\ -3 & 2 \end{bmatrix}$$

$$A^{-1} = \text{Adj } A / |A| = \frac{-1}{5} \begin{bmatrix} -1 & -1 \\ -3 & 2 \end{bmatrix} = \begin{bmatrix} 1/5 & 1/5 \\ 3/5 & -2/5 \end{bmatrix}$$

$$X = A^{-1} B$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = X = \begin{bmatrix} 1/5 & 1/5 \\ 3/5 & -2/5 \end{bmatrix} \begin{bmatrix} 1 \\ 4 \end{bmatrix} = \begin{bmatrix} \frac{1}{5} + \frac{4}{5} \\ \frac{3}{5} - \frac{8}{5} \end{bmatrix} = \begin{bmatrix} \frac{5}{5} \\ -\frac{5}{5} \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$X = 1 \quad y = -1$$

### Finding Inverse of a Matrix Using Matrix Inversion Method

A is matrix. We find the inverse of A ( $A^{-1}$ ) as follows

$$A * (A^{-1}) = I$$

Procedure for finding inverse of a matrix:-

$$A^{-1} = \frac{\text{adj}(A)}{|A|}$$

Adj(A):-

1. Find the minor of each element
2. Find the cofactor of each element
3. Form the cofactor matrix Co(A)
4. Form the adj(A)=[co(A)]<sup>T</sup>

$$A = \begin{bmatrix} 1 & 1 & 2 \\ 5 & 2 & 7 \\ 0 & 1 & 3 \end{bmatrix} \text{ order of } A = 3 \times 3$$

$$|A| = -3$$

$$\text{Minor of } 1 = -1 \quad \text{co-factor of } 1 = -1$$

$$\text{Minor of } 1 = 15 \quad \text{co-factor of } 1 = -15$$

$$\text{Minor of } 2 = 5 \quad \text{co-factor of } 2 = 5$$

Minor of 5= 1      co-factor of 5= -1  
 Minor of 2= 3      co-factor of 2= 3  
 Minor of 7= 1      co-factor of 7= -1  
 Minor of 0= 3      co-factor of 0 = 3  
 Minor of 1= -3      co-factor of 1= 3  
 Minor of 3= -3      co-factor of 3= -3

Co-factor matrix:-

$$\text{Co}(A) = \begin{bmatrix} -1 & -15 & 5 \\ -1 & 3 & -1 \\ 3 & 3 & -3 \end{bmatrix} \quad \text{Adj}(A) = \begin{bmatrix} -1 & -1 & 3 \\ -15 & 3 & 3 \\ 5 & -1 & -3 \end{bmatrix}$$

$$A^{-1} = \frac{\text{adj}(A)}{|A|} = \frac{1}{-3} \begin{bmatrix} -1 & -1 & 3 \\ -15 & 3 & 3 \\ 5 & -1 & -3 \end{bmatrix}$$

$$A = \begin{vmatrix} 1 & -2 \\ 4 & 3 \end{vmatrix}$$

$$M_{11}=3 \quad C_{11}= (-1)^2 \times 3 = 3 \quad \text{ADJ} \quad A$$

$$= \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix}^T$$

$$M_{12}=4 \quad C_{12}= (-1)^3 \times 4 = -4 \quad \text{ADJ } A = \begin{vmatrix} C_{11} & C_{21} \\ C_{12} & C_{22} \end{vmatrix}$$

$$M_{21}=-2 \quad C_{21}= (-1)^3 \times -2 = 2 \quad \text{ADJ } A = \begin{vmatrix} 3 & 2 \\ -4 & 1 \end{vmatrix}$$

$$M_{22}=1 \quad C_{22}= (-1)^4 \times 1 = 1$$

## TRIGNNOMETRY

### ASTC RULE

$$\sin(90-\theta)=\cos\theta$$

$$\cos(90-\theta)=\sin\theta$$

$$\tan(90-\theta)=\cot\theta$$

$$\cot(90-\theta)=\tan\theta$$

$$\sec(90-\theta)=\operatorname{cosec}\theta$$

$$\operatorname{cosec}(90-\theta)=\sec\theta$$

2<sup>nd</sup> quadrant (90°-180°)  
Sin, cosec +ve  
Tan, sec, cos, cot -ve

1<sup>st</sup> quadrant (0°-90°)  
Sin, cos, tan, sec, cosec, cot  
All +ve

3<sup>rd</sup> quadrant (180°-270°)  
Tan, cot +ve  
Sin, sec, cos, cosec -ve

4<sup>th</sup> quadrant (270°-360°)  
Cos, sec +ve  
Sin, cosec, cot, tan -ve

$\Theta$ ( $0 < \theta < 90$ )	$90-\theta$ (1 <sup>st</sup> qrd.)	$90+\theta$ (2 <sup>nd</sup> qrd.)	$180-\theta$ (2 <sup>nd</sup> qrd.)	$180+\theta$ (3 <sup>rd</sup> qrd.)
Sin	cos $\theta$	cos $\theta$	sin $\theta$	-sin $\theta$
Cos	sin $\theta$	-sin $\theta$	-Cos $\theta$	-Cos $\theta$
Tan	cot $\theta$	-cot $\theta$	-Tan $\theta$	Tan $\theta$
Cot	tan $\theta$	-tan $\theta$	-Cot $\theta$	Cot $\theta$
sec	cosec $\theta$	-cosec $\theta$	-Sec $\theta$	-Sec $\theta$
cosec	sec $\theta$	sec $\theta$	cosec $\theta$	-cosec $\theta$

**2275 = 90 X 25 + 25 2<sup>ND</sup> QUADRENT**

**2275 = 360 X 6 + 115**

**Sin 3754° = sin (90 X 41 + 64) = cos 64 2<sup>ND</sup> QUADRENT**

$$3754 = 360 \times 10 + 154$$

$$3333 = 360 \times 9 + 93$$


$$3333 = 90 \times 37 + 3$$

$$\sin 3754^\circ = \sin 154^\circ = \sin(90+64) = \cos 64 = +ve$$

Trigonometric Ratios:-

$$\sin \theta = \frac{p}{h}; \cos \theta = \frac{b}{h}; \tan \theta = \frac{p}{b}; \cot \theta = \frac{b}{p}$$

$$\sec \theta = \frac{h}{b}; \operatorname{cosec} \theta = \frac{h}{p}$$



Pythagoras theorem

$$p^2 + b^2 = h^2 ; p\text{- Perpendicular, } b\text{- Base, } h\text{- Hypotenuse}$$

$$\sin^2 \theta + \cos^2 \theta = \left(\frac{p}{h}\right)^2 + \left(\frac{b}{h}\right)^2 = \frac{p^2}{h^2} + \frac{b^2}{h^2} = \frac{p^2 + b^2}{h^2} = \frac{h^2}{h^2} = 1$$

$$\sec^2 \theta - \tan^2 \theta = 1$$

$$\operatorname{cosec}^2 \theta - \cot^2 \theta = 1$$

$$\sec \theta = 1/\cos \theta$$

$$\cot \theta = 1/\tan \theta$$

$$\operatorname{cosec} \theta = 1/\sin \theta$$

$\Theta$ ( $0 < \theta < 90$ )	$90 - \theta$ (1 <sup>st</sup> qrd.)	$90 + \theta$ (2 <sup>nd</sup> qrd.)	$180 - \theta$ (2 <sup>nd</sup> qrd.)	$180 + \theta$ (3 <sup>rd</sup> qrd.)
Sin	cos $\theta$	cos $\theta$	sin $\theta$	-sin $\theta$
Cos	sin $\theta$	-sin $\theta$	-Cos $\theta$	-Cos $\theta$
Tan	cot $\theta$	-cot $\theta$	-Tan $\theta$	Tan $\theta$
Cot	tan $\theta$	-tan $\theta$	-Cot $\theta$	Cot $\theta$
sec	cosec $\theta$	-cosec $\theta$	-Sec $\theta$	-Sec $\theta$
cosec	sec $\theta$	sec $\theta$	cosec $\theta$	-cosec $\theta$

$F(-x) = -f(x)$  odd function

$f(-x) = f(x)$  even function

Cos, sec even function

Sin, tan, cosec, cot odd function

$$\sin(-3333^\circ) = -\sin 3333^\circ = -\sin(90^\circ \times 37 + 3^\circ) = -\cos 3^\circ$$

$$\sin(3423^\circ) = \sin(90^\circ \times 38 + 3^\circ) = -\sin 3^\circ$$

$$\cot(3425^\circ) = \cot(90^\circ \times 38 + 5^\circ) = +\cot 5^\circ$$

$$\operatorname{cosec}(6255^\circ) = \operatorname{cosec}(90^\circ \times 69 + 45^\circ) = +\sec 45^\circ = \sqrt{2}$$

$$\sec(-7634^\circ) = \sec 7634^\circ = \sec(90^\circ \times 84 + 74^\circ) = +\sec 74^\circ$$

1. Find the value of  $\cos 1^\circ \cdot \cos 2^\circ \cdot \dots \cdot \cos 100^\circ$ .

Solution:-

$$\begin{aligned} & \cos 1^\circ \cdot \cos 2^\circ \cdot \dots \cdot \cos 100^\circ \\ &= \cos 1^\circ \cdot \cos 2^\circ \cdot \dots \cdot \cos 90^\circ \cdot \dots \cdot \cos 100^\circ \\ &= 0 \text{ [as } \cos 90^\circ = 0\text{]} \end{aligned}$$

find the value of  $\tan 1^\circ \cdot \tan 2^\circ \cdot \dots \cdot \tan 89^\circ$

$$\begin{aligned} & \tan 1^\circ \cdot \tan 2^\circ \cdot \dots \cdot \tan 89^\circ \\ &= \tan 1^\circ \cdot \tan 2^\circ \cdot \dots \cdot \tan 44^\circ \cdot \tan 45^\circ \cdot \tan 46^\circ \cdot \dots \cdot \tan 88^\circ \cdot \tan 89^\circ \\ &= \tan 1^\circ \cdot \tan 2^\circ \cdot \dots \cdot \tan 44^\circ \cdot 1 \cdot \tan(90^\circ - 44^\circ) \cdot \dots \cdot \tan(90^\circ - 2^\circ) \cdot \tan(90^\circ - 1^\circ) \\ &= \tan 1^\circ \cdot \tan 2^\circ \cdot \dots \cdot \tan 44^\circ \cdot 1 \cdot \cot 44^\circ \cdot \dots \cdot \cot 2^\circ \cdot \cot 1^\circ \\ &= (\tan 1^\circ \cdot \cot 1^\circ) \cdot (\tan 2^\circ \cdot \cot 2^\circ) \cdot \dots \cdot (\tan 44^\circ \cdot \cot 44^\circ) \cdot 1 \\ &= 1 \cdot 1 \cdot 1 \cdot \dots \cdot 1 \\ &= 1 \end{aligned}$$

2. find the value of  $\cos 24^\circ + \cos 5^\circ + \cos 175^\circ + \cos 204^\circ + \cos 300^\circ$

**solution:**  $\cos 24^\circ + \cos 5^\circ + \cos 175^\circ + \cos 204^\circ + \cos 300^\circ$

$$\begin{aligned} &= \cos 24^\circ + \cos 5^\circ + \cos(180^\circ - 5^\circ) + \cos(180^\circ + 24^\circ) + \cos(90^\circ \times 3 + 30^\circ) \\ &= \cos 24^\circ + \cos 5^\circ - \cos 5^\circ - \cos 24^\circ + \sin 30^\circ \end{aligned}$$

$$= \sin 30^\circ = 1/2$$

$$\text{Sol} = \cos 24^\circ + \cos 5^\circ + \cos (180^\circ - 5^\circ) + \cos (180^\circ + 24^\circ) + \cos (360^\circ - 60^\circ)$$

$$= \cos 24^\circ + \cos 5^\circ - \cos 5^\circ - \cos 24^\circ + \cos 60^\circ$$

$$= 0 + 0 + 1/2$$

$$= 1/2$$

find the value of  $\tan 1^\circ \cdot \tan 2^\circ \cdot \dots \cdot \tan 100^\circ$

Sol:-

$$\tan 1^\circ \cdot \tan 2^\circ \cdot \dots \cdot \tan 100^\circ$$

$$= \tan 1^\circ \cdot \tan 2^\circ \cdot \dots \cdot \tan 90^\circ \cdot \dots \cdot \tan 100^\circ$$

$$= \infty$$

find the value of  $\begin{vmatrix} \cos x & \sin x \\ \sin x & \cos x \end{vmatrix}$

$$\begin{vmatrix} \cos x & \sin x \\ \sin x & \cos x \end{vmatrix} = \cos^2 x - \sin^2 x = \cos 2x =$$

maximum value 1, minimum value - 1

Some standard formula:-

1.  $\sin(A+B) = \sin A \cdot \cos B + \cos A \cdot \sin B$
2.  $\sin(A-B) = \sin A \cdot \cos B - \cos A \cdot \sin B$
3.  $\cos(A+B) = \cos A \cdot \cos B - \sin A \cdot \sin B$
4.  $\cos(A-B) = \cos A \cdot \cos B + \sin A \cdot \sin B$
5.  $\sin 2A = \sin(A+A) = \sin A \cdot \cos A + \cos A \cdot \sin A = 2 \sin A \cdot \cos A$
6.  $\cos 2A = \cos(A+A) = \cos A \cdot \cos A - \sin A \cdot \sin A = \cos^2 A - \sin^2 A$
7.  $\sin 3A = 3 \sin A - 4 \sin^3 A$

$$\begin{aligned} \sin 3A &= \sin(A+2A) = \sin A \cdot \cos 2A + \cos A \cdot \sin 2A \\ &= \sin A \cdot (\cos^2 A - \sin^2 A) + \cos A \cdot 2 \sin A \cdot \cos A \\ &= \sin A (1 - \sin^2 A - \sin^2 A) + 2 \sin A \cdot \cos^2 A \\ &= \sin A - 2 \sin^3 A + 2 \sin A (1 - \sin^2 A) \\ &= \sin A - 2 \sin^3 A + 2 \sin A - 2 \sin^3 A \\ &= 3 \sin A - 4 \sin^3 A \end{aligned}$$



$$8. \cos 3A = \cos(A+2A) = 4\cos^3 A - 3\cos A$$

**Problem-  $\sin 18^\circ$**

$$\Theta = 18^\circ$$

$$\Rightarrow 5\theta = 90^\circ$$

$$\Rightarrow 3\theta + 2\theta = 90^\circ$$

$$\Rightarrow 2\theta = 90^\circ - 3\theta$$

$$\Rightarrow \sin 2\theta = \sin(90^\circ - 3\theta)$$

$$\Rightarrow \sin 2\theta = \cos 3\theta$$

$$\Rightarrow 2\sin\theta \cdot \cos\theta = 4\cos^3\theta - 3\cos\theta$$

$$\Rightarrow 2\sin\theta \cdot \cos\theta = \cos\theta(4\cos^2\theta - 3)$$

$$\Rightarrow 2\sin\theta = (4\cos^2\theta - 3)$$

$$\Rightarrow 2\sin\theta = 4(1 - \sin^2\theta) - 3 = 4 - 4\sin^2\theta - 3 = 1 - 4\sin^2\theta$$

$$\Rightarrow 2\sin\theta = 1 - 4\sin^2\theta$$

$$\Rightarrow 2\sin\theta + 4\sin^2\theta - 1 = 0$$

$$\Rightarrow 4\sin^2\theta + 2\sin\theta - 1 = 0$$

$$Ax^2 + bx + c = 0 \quad x = \frac{-b \pm \sqrt{b^2 - 4a \cdot c}}{2a}$$

$$A=4, b=2, c=-1$$

$$\sin\theta = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 4 \cdot (-1)}}{2 \cdot 4} = \frac{-2 \pm \sqrt{20}}{2 \cdot 4} = \frac{-2 \pm 2\sqrt{5}}{2 \cdot 4} = \frac{-1 \pm \sqrt{5}}{4}$$

$$\sin 18^\circ = \frac{-1 + \sqrt{5}}{4}$$

$$\text{Therefore } \sin 18^\circ = \frac{-1 + \sqrt{5}}{4} \quad (\sin 18 \text{ is always +ve})$$

$$\cos 18^\circ, \sin 36^\circ, \cos 36^\circ$$

If  $\sin\theta + \operatorname{cosec}\theta = 2$ . how that  $\sin^n\theta + \operatorname{cosec}^n\theta = 2$  for all positive integers  $n$ .

$$\sin\theta + \operatorname{cosec}\theta = 2$$

$$\sin\theta + \frac{1}{\sin\theta} = 2$$

$$\sin^2\theta + 1 = 2\sin\theta$$

$$\sin^2\theta - 2\sin\theta + 1 = 0$$

$$\sin^2\theta - 2 \cdot 1 \cdot \sin\theta + 1 = 0$$

$$(\sin\theta - 1)^2 = 0$$

$$\sin\theta = 1, \operatorname{cosec}\theta = 1$$

$$\sin^n\theta + \operatorname{cosec}^n\theta = 1^n + 1^n = 1 + 1 = 2$$

**Find the maximum value of  $3\sin x + 4\cos x$  ?**

$$3\sin x + 4\cos x$$

$$\text{Lect } 3 = r \cos\theta, 4 = r \sin\theta$$

$$r \cos\theta \sin x + r \sin\theta \cos x$$

$$r(\sin x \cdot \cos\theta + \cos x \cdot \sin\theta)$$

$$3\cos x + 4\sin x$$

$$\text{Lect } 3 = r \sin\theta, 4 = r \cos\theta$$

$$r \sin\theta \cos x + r \cos\theta \sin x$$

$$r(\sin(x+\theta))$$

$$3^2 + 4^2 = r^2 \cos^2 \theta + r^2 \sin^2 \theta$$

$$9+16=r^2(\cos^2 \theta + \sin^2 \theta) = r^2 \cdot 1$$

$$25 = r^2$$

$$r = +5, -5$$

maximum value of  $3 \sin x + 4 \cos x$  is 5

minimum value of  $3 \sin x + 4 \cos x$  is -5

1.  $\sin(A+B) = \sin A \cdot \cos B + \cos A \cdot \sin B$

2.  $\sin(A-B) = \sin A \cdot \cos B - \cos A \cdot \sin B$

3.  $\cos(A+B) = \cos A \cdot \cos B - \sin A \cdot \sin B$

4.  $\cos(A-B) = \cos A \cdot \cos B + \sin A \cdot \sin B$

5.  $\sin 2A = \sin(A+A) = \sin A \cdot \cos A + \cos A \cdot \sin A = 2 \sin A \cdot \cos A$

6.  $\cos 2A = \cos(A+A) = \cos A \cdot \cos A - \sin A \cdot \sin A = \cos^2 A - \sin^2 A$

7.  $\sin 3A = 3 \sin A - 4 \sin^3 A$

8.  $\sin 2A = \frac{2 \tan A}{1 + \tan^2 A}$

$$\sin A = \frac{2 \tan \frac{A}{2}}{1 + \tan^2 \frac{A}{2}}$$

9.  $\cos 2A = \frac{1 - \tan^2 A}{1 + \tan^2 A}$

$$\cos A = \frac{1 - \tan^2 \frac{A}{2}}{1 + \tan^2 \frac{A}{2}}$$

10.  $\cos 2A = \cos^2 A - \sin^2 A = 1 - \sin^2 A - \sin^2 A = 1 - 2 \sin^2 A$   
 $1 - 2 \sin^2 A = \cos 2A$

$$1 - \cos 2A = 2 \sin^2 A$$

$$1 + \cos 2A = 2 \cos^2 A$$

$$\sin^2 A = \frac{1 - \cos 2A}{2}$$

11.  $\sin A = \sqrt{\frac{1 - \cos 2A}{2}}$

$$\sin \frac{A}{2} = \sqrt{\frac{1 - \cos A}{2}}$$

12.  $\cos A = \sqrt{\frac{1 + \cos 2A}{2}}$

$$\cos \frac{A}{2} = \sqrt{\frac{1 + \cos A}{2}}$$

13.  $\tan A = \sqrt{\frac{1 - \cos 2A}{1 + \cos 2A}}$

$$\tan \frac{A}{2} = \sqrt{\frac{1 - \cos A}{1 + \cos A}}$$

14.  $\sin C + \sin D = 2 \sin \frac{C+D}{2} \cdot \cos \frac{C-D}{2}$

15.  $\sin C - \sin D = 2 \cos \frac{C+D}{2} \cdot \sin \frac{C-D}{2}$

16.  $\cos C + \cos D = 2 \cos \frac{C+D}{2} \cdot \cos \frac{C-D}{2}$

17.  $\cos C - \cos D = 2 \sin \frac{C+D}{2} \cdot \sin \frac{D-C}{2}$

**PROBLEM, :-** If  $A+B+C=\pi$  then prove that(PROVE THAT IN A TRIANGLE)

$$\sin 2A + \sin 2B + \sin 2C = 4 \sin A \cdot \sin B \cdot \sin C$$

Proof:-

L.H.S

$$\begin{aligned} & \sin 2A + \sin 2B + \sin 2C \\ &= 2 \sin \frac{2A+2B}{2} \cdot \cos \frac{2A-2B}{2} + 2 \sin C \cdot \cos C \\ &= 2 \sin (A+B) \cdot \cos(A-B) + 2 \sin C \cdot \cos C \\ & \text{[given } A+B+C=\pi \Rightarrow A+B=\pi-C \text{]} \\ &= 2 \sin(\pi-C) \cdot (\cos A \cdot \cos B + \sin A \cdot \sin B) + 2 \sin C \cdot \cos C \\ &= 2 \sin C \cdot (\cos A \cdot \cos B + \sin A \cdot \sin B) + 2 \sin C \cdot \cos C \\ &= 2 \sin C \cdot \cos A \cdot \cos B + 2 \sin C \cdot \sin A \cdot \sin B + 2 \sin C \cdot \cos C \\ &= 2 \sin C \cdot \cos A \cdot \cos B + 2 \sin C \cdot \cos C + 2 \sin A \cdot \sin B \cdot \sin C \\ &= 2 \sin C [\cos A \cdot \cos B + \cos C] + 2 \sin A \cdot \sin B \cdot \sin C \\ &= 2 \sin C [\cos A \cdot \cos B + \cos(\pi - (A+B))] + 2 \sin A \cdot \sin B \cdot \sin C \\ &= 2 \sin C [\cos A \cdot \cos B - \cos(A+B)] + 2 \sin A \cdot \sin B \cdot \sin C \\ &= 2 \sin C [\cos A \cdot \cos B - (\cos A \cdot \cos B - \sin A \cdot \sin B)] + 2 \sin A \cdot \sin B \cdot \sin C \\ &= 2 \sin C [\cos A \cdot \cos B - \cos A \cdot \cos B + \sin A \cdot \sin B] + 2 \sin A \cdot \sin B \cdot \sin C \\ &= 2 \sin C \cdot \sin A \cdot \sin B + 2 \sin A \cdot \sin B \cdot \sin C \\ &= 2 \sin A \cdot \sin B \cdot \sin C + 2 \sin A \cdot \sin B \cdot \sin C \\ &= 4 \sin A \cdot \sin B \cdot \sin C \end{aligned}$$

**h.w-PROBLEM, :-** If  $A+B+C=\pi$  then prove that(PROVE THAT IN A TRIANGLE)

$$\cos 2A + \cos 2B + \cos 2C + 1 = -4 \cos A \cdot \cos B \cdot \cos C$$

Show that the equation  $\sin \theta = a + \frac{1}{a}$  does not have a solution for every real number  $a \neq 0$

$$\sin \theta = \frac{a^2 + 1}{a}$$

If  $a=0$

Then  $\sin\theta = 1$  we have solution for this .

Suppose  $a \neq 0$

For any positive value  $a \frac{a^2+1}{a} > 1$

For any negative value  $a \frac{a^2+1}{a} < -1$

But we know range of  $\sin\theta$  is  $[-1, 1]$  so for any value

$\sin\theta = a + \frac{1}{a}$  does not have a solution for every real number  $a \neq 0$

**Find the maximum value of  $5\sin x + 12 \cos x$**

Lect  $5 = r \cos\theta$  ,  $12 = r \sin\theta$

$r \cos\theta \sin x + r \sin\theta \cos x$

$r(\sin x \cdot \cos\theta + \cos x \cdot \sin\theta)$

$r(\sin(x+\theta))$

$5^2 + 12^2 = r^2 \cos^2\theta + r^2 \sin^2\theta$

$25+144 = r^2(\cos^2\theta + \sin^2\theta) = r^2 \cdot 1$

$169 = r^2$

$r = 13, -13$

**Find the maximum value of  $5\sin x + 12 \cos x$  is 13**

$2+3\sin x+4\cos x$

$3 \sin x + 4 \cos x$

Lect  $3 = r \cos\theta$  ,  $4 = r \sin\theta$

$2+r \cos\theta \sin x + r \sin\theta \cos x$

$2+r(\sin x \cdot \cos\theta + \cos x \cdot \sin\theta)$

$2+r(\sin(x+\theta))$

$3^2 + 4^2 = r^2 \cos^2\theta + r^2 \sin^2\theta$

$9+16 = r^2(\cos^2\theta + \sin^2\theta) = r^2 \cdot 1$

$25 = r^2$

$R = 5, -5$

Maximum value =  $2+5 = 7$

Minimum value =  $2-5 = -3$

**Find the maximum value of  $3\sin x+4\cos x-2$**

$3\sin x+4\cos x-2$

$3 \sin x + 4 \cos x - 2$

Lect  $3 = r \cos\theta$  ,  $4 = r \sin\theta$

$r \cos\theta \sin x + r \sin\theta \cos x - 2$

$r(\sin x \cdot \cos\theta + \cos x \cdot \sin\theta) - 2$

$r(\sin(x+\theta)) - 2$

$3^2 + 4^2 = r^2 \cos^2\theta + r^2 \sin^2\theta$

$$9+16=r^2(\cos^2 \theta + \sin^2 \theta) = r^2 \cdot 1$$

$$25 = r^2$$

$$R = 5, -5$$

$$\text{Maximum value} = 5 - 2 = 3$$

$$\text{Minimum value} = -5 - 2 = -7$$

Find the maximum value of  $2 - 3\sin x - 4\cos x$

$$2 - 3\sin x - 4\cos x$$

$$2 - 3\sin x - 4\cos x$$

$$\text{Let } 3 = r \cos \theta, 4 = r \sin \theta$$

$$2 - r \cos \theta \sin x - r \sin \theta \cos x$$

$$2 - r(\sin x \cdot \cos \theta + \cos x \cdot \sin \theta)$$

$$2 - r(\sin(x + \theta))$$

$$3^2 + 4^2 = r^2 \cos^2 \theta + r^2 \sin^2 \theta$$

$$9 + 16 = r^2(\cos^2 \theta + \sin^2 \theta) = r^2 \cdot 1$$

$$25 = r^2$$

$$R = 5, -5$$


$$\text{Minimum value} = 2 - 5 = -3$$

$$\text{Maximum value} = 2 - (-5) = 7$$

### INVERSE TRIGNOMETRIC FUNCTION:

$F: A \rightarrow B$  then  $f^{-1}: B \rightarrow A$

$\sin: \mathbb{R} \rightarrow [-1, 1]$      $\sin^{-1}: [-1, 1] \rightarrow \mathbb{R}$



$$\sin(\sin^{-1}x) = x \quad \sin^{-1}(\sin x) = x \quad \sin^{-1}(1) = \sin^{-1}(\sin 90) = 90^\circ$$

$$\cos(\cos^{-1}x) = x \quad \cos^{-1}(\cos x) = x$$

$$\tan(\tan^{-1}x) = x \quad \tan^{-1}(\tan x) = x \quad \tan^{-1}\left(\frac{1}{\sqrt{3}}\right) = \tan^{-1}(\tan 30) = 30^\circ$$

$$\sec(\sec^{-1}x) = x \quad \sec^{-1}(\sec x) = x \quad \sec^{-1}(2) = \sec^{-1}(\sec 60) = 60^\circ$$

$$\operatorname{cosec}(\operatorname{cosec}^{-1}x) = x \quad \operatorname{cosec}^{-1}(\operatorname{cosec} x) = x$$

$$\cot(\cot^{-1}x) = x \quad \cot^{-1}(\cot x) = x$$

$$\operatorname{cosec}^{-1}x = \sin^{-1}\left(\frac{1}{x}\right); |x| \geq 1, \quad \sin^{-1}\left(\frac{1}{x}\right) = \operatorname{cosec}^{-1}(x)$$

$$\sec^{-1}x = \cos^{-1}\left(\frac{1}{x}\right); |x| \geq 1, \quad \sec^{-1}\left(\frac{1}{x}\right) = \cos^{-1}(x)$$

$$\cot^{-1}x = \tan^{-1}\left(\frac{1}{x}\right), x > 0 \quad \cot^{-1}\left(\frac{1}{x}\right) = \tan^{-1}(x), x > 0$$

$$\cot^{-1}x = \pi + \tan^{-1}\left(\frac{1}{x}\right), x < 0 \quad \cot^{-1}\left(\frac{1}{x}\right) = \pi + \tan^{-1}(x), x < 0$$

$$\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}, x \in [-1, 1]$$

$$\tan^{-1}x + \cot^{-1}x = \frac{\pi}{2}, x \in \mathbb{R}$$

$$\sec^{-1}x + \operatorname{cosec}^{-1}x = \frac{\pi}{2}, x \in (-\infty, -1] \cup [1, \infty)$$

$$\tan^{-1}x + \tan^{-1}y = \begin{cases} \tan^{-1}\frac{x+y}{1-xy} & \text{if } xy < 1 \\ \pi + \tan^{-1}\frac{x+y}{1-xy} & \text{if } xy > 1, x > 0, y > 0 \\ -\pi + \tan^{-1}\frac{x+y}{1-xy} & \text{if } xy > 1, x < 0, y < 0 \end{cases}$$

$$\tan^{-1}x - \tan^{-1}y = \tan^{-1}\frac{x-y}{1+xy} \text{ if } x \geq 0, y \geq 0$$

$$\text{problem 1 : } \tan^{-1}\left(\frac{1}{2}\right) + \tan^{-1}\left(\frac{1}{3}\right)$$

$$[x=1/2, y=1/3 \Rightarrow xy=1/6 < 1]$$

$$= \tan^{-1} \frac{\left(\frac{1}{2}\right) + \left(\frac{1}{3}\right)}{1 - \left(\frac{1}{2}\right)\left(\frac{1}{3}\right)} = \tan^{-1} \frac{\left(\frac{5}{6}\right)}{\left(\frac{5}{6}\right)} = \tan^{-1} 1 = \tan^{-1}(\tan \frac{\pi}{4}) = \pi/4$$

$$\text{Problem 2: Find the value of } \cos \tan^{-1} \cot \cos^{-1} \left(\frac{\sqrt{3}}{2}\right)?$$

$$\cos \tan^{-1} \cot \cos^{-1} \left(\frac{\sqrt{3}}{2}\right)$$

$$= \cos \tan^{-1} \cot \cos^{-1} \left(\cos \frac{\pi}{6}\right)$$

$$= \cos \tan^{-1} \cot 30^\circ$$

$$= \cos \tan^{-1} \sqrt{3}$$

$$= \cos \tan^{-1} (\tan 60^\circ)$$

$$= \cos 60^\circ$$

$$= 1/2$$

$$\text{q- } \sin^{-1} \frac{1}{\sqrt{5}} + \cos^{-1} \frac{3}{\sqrt{10}} = ?$$

$$\sin^{-1} x + \cos^{-1} y = \sin^{-1} x + \left(\frac{\pi}{2} - \sin^{-1} y\right)$$

$$= \frac{\pi}{2} + \sin^{-1} x - \sin^{-1} y = \frac{\pi}{2} + \sin^{-1}(x\sqrt{1-y^2} - y\sqrt{1-x^2})$$

$$\sin^{-1} x + \cos^{-1} y = \frac{\pi}{2} + \sin^{-1}(x\sqrt{1-y^2} - y\sqrt{1-x^2})$$

$$\sin^{-1} \frac{1}{\sqrt{5}} + \cos^{-1} \frac{3}{\sqrt{10}} = \frac{\pi}{2} + \sin^{-1} \left( \frac{1}{\sqrt{5}} \sqrt{1 - \frac{3^2}{10}} - \frac{3}{\sqrt{10}} \sqrt{1 - \frac{1^2}{5}} \right)$$

$$= \frac{\pi}{2} + \sin^{-1} \left( \frac{1}{\sqrt{5}} \sqrt{1 - \frac{9}{10}} - \frac{3}{\sqrt{10}} \sqrt{1 - \frac{1}{5}} \right)$$

$$= \frac{\pi}{2} + \sin^{-1} \left( \frac{1}{\sqrt{5}} * \frac{1}{\sqrt{10}} - \frac{3}{\sqrt{10}} * \frac{2}{\sqrt{5}} \right) = \frac{\pi}{2} + \sin^{-1} \left( -\frac{5}{5\sqrt{2}} \right)$$

$$= \frac{\pi}{2} + \sin^{-1}\left(-\frac{1}{\sqrt{2}}\right) = \frac{\pi}{2} + \sin^{-1}\left(\sin\left(-\frac{\pi}{4}\right)\right) = \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}$$

q-4  $\cos^{-1} x + \sin^{-1} x = \pi$  what is the value of x

$$4 \cos^{-1} x + \sin^{-1} x = \pi$$

$$2.2 \cos^{-1} x + \sin^{-1} x = \pi$$

$$2 \cdot \cos^{-1}(2x^2 - 1) + \sin^{-1} x = \pi$$

$$\cos^{-1}(2(2x^2 - 1)^2 - 1) + \sin^{-1} x = \pi$$

$$\cos^{-1}(2(2x^2 - 1)^2 - 1) + \frac{\pi}{2} - \cos^{-1} x = \pi$$

$$\cos^{-1}(2(2x^2 - 1)^2 - 1) - \cos^{-1} x = \pi - \frac{\pi}{2} = \frac{\pi}{2}$$

find  $\sec^2(\tan^{-1} 2) + \operatorname{cosec}^2(\cot^{-1} 3)$

$$\sec^2(\tan^{-1} 2) + \operatorname{cosec}^2(\cot^{-1} 3)$$

$$= 1 + \tan^2(\tan^{-1} 2) + 1 + \cot^2(\cot^{-1} 3)$$

$$= 1 + (\tan(\tan^{-1} 2))^2 + 1 + (\cot(\cot^{-1} 3))^2$$

$$= 1 + 2^2 + 3^2 = 1 + 4 + 1 + 9 = 15$$

$$\tan^2 x = (\tan x)^2$$

If  $\tan^{-1} 2$  and  $\tan^{-1} 3$  are two angle of a triangle then third angle=?

$$\tan^{-1} 2 + \tan^{-1} 3 + \tan^{-1} x = 180^\circ$$

$$\tan^{-1}\left(\frac{2+3+x-6x}{5-5x}\right) = 180^\circ$$

$$2 + 3 + x - 6x = 0$$

$$5 - 5x = 0, x = 1$$


$$\tan^{-1} 1 = 45^\circ$$



## Co-ordinate geometry in two dimensions

### Cartesian product:-

A & B are two sets then their  $A \times B$  is the order pair  $(x,y)$  such that  $x \in A$  &  $y \in B$ .  
 $A \times B = \{(x,y) : x \in A, y \in B\}$ .



### Distance formula:-

$$(x_1, y_1) \text{-----} (x_2, y_2)$$

Let  $P(x_1, y_1)$  &  $Q(x_2, y_2)$  are two points on the co-ordinate geometry. Then their distance can be calculated as follows

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

ex:- distance between (3,5) & (2,1)

$$x_1 = 3, y_1 = 5, x_2 = 2, y_2 = 1$$

$$d = \sqrt{(2 - 3)^2 + (1 - 5)^2} = \sqrt{(-1)^2 + (-4)^2} = \sqrt{1 + 16} = \sqrt{17}$$

ex:- find the distance between two points (-1,1) and (3,3);

$$d = \sqrt{(3 - (-1))^2 + (3 - 1)^2} = \sqrt{(4)^2 + (2)^2} = \sqrt{16 + 4} = \sqrt{20}$$

:- find the distance between two points (2,0) and (-3,7)

$$d = \sqrt{(-3 - 2)^2 + (7 - 0)^2} = \sqrt{(-5)^2 + (7)^2} = \sqrt{25 + 49} = \sqrt{74}$$

### **Division Formula (Internal):-**

Suppose  $P(x,y)$  intersects the line segment joining the points  $(x_1, y_1)$  &  $(x_2, y_2)$  internally in the ration  $m:n$  then the co-ordinate of  $(x,y)$  is

$$x = \frac{mx_2 + nx_1}{m+n} \text{ and } y = \frac{my_2 + ny_1}{m+n}$$

Ex- find the point which intersects the line segment joining the points (5,4) and (2,1) internally in the ration 2:1?

Sol- here  $x_1 = 5$   $x_2 = 2$   $y_1 = 4$   $y_2 = 1$   $m = 2$ ,  $n = 1$

Therefore by internal division formula

The point  $P(x,y)$  which intersects the line segment joining the points (3,4) and (2,1) is

$$x = \frac{mx_2 + nx_1}{m+n} \text{ and } y = \frac{my_2 + ny_1}{m+n}$$

$$x = \frac{2 \cdot 2 + 1 \cdot 5}{2+1}; y = \frac{2 \cdot 1 + 1 \cdot 4}{2+1}$$

$$x = 9/3; y = 6/3 = 2$$

Therefore the point is (3,2)

### **Division Formula (External):-**

Suppose  $P(x,y)$  intersects the line segment joining the points  $(x_1, y_1)$  &  $(x_2, y_2)$  externally in the ration  $m:n$  then the co-ordinate of  $(x,y)$  is

$$x = \frac{mx_2 - nx_1}{m-n} \text{ and } y = \frac{my_2 - ny_1}{m-n}$$

Ex- find the point which intersects the line segment joining the points (3,4) and (2,1) externally in the ration 2:1

here  $x_1 = 3$   $x_2 = 2$   $y_1 = 4$   $y_2 = 1$   $m = 2$ ,  $n = 1$

$$x = \frac{2 \cdot 2 - 1 \cdot 3}{2 - 1} \text{ and } y = \frac{2 \cdot 1 - 1 \cdot 4}{2 - 1}$$

$$x = 1/1 = 1, y = (2 - 4)/1 = -2$$

\*\*\* Suppose  $P(x, y)$  bisects the line segment joining the points  $(x_1, y_1)$  &  $(x_2, y_2)$  then the co-ordinate of  $(x, y)$  is

$$x = \frac{x_2 + x_1}{2} \text{ and } y = \frac{y_2 + y_1}{2}$$

### Area of Triangle:-

Suppose ABC is a triangle with A  $(x_1, y_1)$ ; B  $(x_2, y_2)$  & C  $(x_3, y_3)$  are the vertices. Then the area of the triangle ABC is

$$\Delta = \frac{1}{2} \begin{vmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{vmatrix}$$

Ex:- find the area of the triangle whose vertices are  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$

Solution- Let ABC is the triangle with vertices as A $(0, 1)$ , B $(1, 0)$  and C $(1, 1)$  then the its area is

$$\Delta = \frac{1}{2} \begin{vmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{vmatrix} = \frac{1}{2} [1(1-0) - 1(0-1) + 1(0-1)] = \frac{1}{2} (1+1-1) = \frac{1}{2} \text{ sq. unit}$$

**If the area of triangle ABC is zero then the points A  $(x_1, y_1)$ ; B  $(x_2, y_2)$  & C  $(x_3, y_3)$  are collinear.**


1.h.w= Find the area of the triangle whose vertices are  $(0, 0)$ ,  $(2, 0)$  and  $(5, 0)$

2.find the mid point of line joining  $(2, 3)$  and  $(4, 5)$

### Slope of a line:-

Suppose a line L makes an angle  $\theta$  with respects to X-axis.

Then the slope of L is the  $\tan\theta$ . Slope of a line is denoted by m. here  $\theta$  is the angle of inclination.



ex:- let angle of inclination of the line L is  $60^\circ$ . Then what is it slope.

Here  $\theta = 60^\circ$  therefore slope,  $m = \tan 60^\circ = \sqrt{3}$

Find the slope of line whose angle of inclination is


- i)  $45^\circ$  ( $m=1$ )    ii)  $30^\circ$  ( $m= 1/\sqrt{3}$ )    iii)  $120^\circ$  ( $m=-\sqrt{3}$ )
- ii) Find the angle of inclination of the line whose slope is

- i)  $1$  ( $\theta=45^\circ$ )    ii)  $\sqrt{3}$  ( $\theta=60^\circ$ )    iii)  $1/\sqrt{3}$  ( $\theta=30^\circ$ )    iv)  $0$  ( $\theta=0^\circ$ )

### Angle between two lines:-

Suppose  $L_1$  &  $L_2$  are two lines with slope  $m_1$  &  $m_2$  respectively. Then the angle between two line  $L_1$  &  $L_2$  is

$$\theta = \tan^{-1}\left(\frac{m_1 - m_2}{1 + m_1 m_2}\right)$$



Ex:- find the angle between two lines whose slopes are 1 & 0.

$$\theta = \tan^{-1}\left(\frac{m_1 - m_2}{1 + m_1 m_2}\right)$$

$$\Theta = \tan^{-1}\left(\frac{1 - 0}{1 + 1 \cdot 0}\right) = \tan^{-1} 1 = \tan^{-1}(\tan 45^\circ) = 45^\circ = \pi/4$$

$$\frac{m_1 - m_2}{1 + m_1 m_2} = 0$$

$$= m_1 - m_2 = 0 \quad m_1 = m_2 \text{ parallel}$$

$$\frac{m_1 - m_2}{1 + m_1 m_2} = \frac{1}{0}$$

$$0 = 1 + m_1 m_2$$

$$m_1 m_2 = -1 \text{ perpendicular}$$

Suppose  $L_1$  &  $L_2$  are two lines with slope  $m_1$  &  $m_2$  respectively then


i)  $L_1 \parallel L_2 \leftrightarrow m_1 = m_2$  (condition of Parallelism)

ii)  $L_1 \perp L_2 \leftrightarrow m_1 m_2 = -1 \Rightarrow m_1 = -\frac{1}{m_2}$  (condition of Perpendicularity)

Ex:- if  $L_1$  &  $L_2$  are two lines with slopes 1 & -1 then what you can tell about  $L_1$  &  $L_2$ .

$$m_1 m_2 = 1 * -1 = -1 \Rightarrow L_1 \perp L_2$$

## Slope of a line which passes through two given points $(x_1, y_1)$ & $(x_2, y_2)$



$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

ex:- suppose a line passes through  $(1,2)$  and  $(2,3)$  then find it's slope.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - 2}{2 - 1} = 1/1 = 1$$

it's angle of inclination is  $45^\circ$

### Equation of a straight line:-

#### 1. Slope intercept form:-

Equation of a line with slope  $m$  and  $y$ -intercept as  $c$  is  $y = mx + c$

$$m = 2 \quad c = 4$$

$$y = 2x + 4$$

ex:- find the equation of a line with slope 1 and  $Y$ -intercept as 3.

$$Y = x + 3$$

#### 2. Slope Point form:-

Equation of a line with slope  $m$  and which passes through  $(x_1, y_1)$  is  $y - y_1 = m(x - x_1)$

Ex:- Find the equation of a line with slope 1 and passes through  $(2,3)$

$$x_1 = 2, y_1 = 3 \quad m = 1$$

$$y - 3 = 1(x - 2)$$




$$y - 3 = x - 2$$

$$y - x - 1 = 0$$

$$y = x + 1$$

#### 3. Two Point form:-

Equation of a line which passes through  $(x_1, y_1)$  &  $(x_2, y_2)$  is



$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1} (x - x_1)$$

Ex:- Find the equation of a line which passes through (-1,2) and (2,3)

$$x_1 = -1, y_1 = 2, x_2 = 2, y_2 = 3$$

$$y - 2 = \frac{3 - 2}{2 - (-1)} (x - (-1))$$

$$y - 2 = \frac{3 - 2}{2 + 1} (x + 1)$$

$$y - 2 = \frac{1}{3} (x + 1)$$

$$3(y - 2) = x + 1$$

$$3y - 6 = x + 1$$

$$3y - x - 7 = 0$$

$$3y = x + 7$$

$$Y = \frac{1}{3} x + \frac{7}{3}$$

#### 4. Intercept form:-

Equation of a line with X-intercept and Y-intercept as a & b respectively is

$$\frac{x}{a} + \frac{y}{b} = 1$$

Ex:- find the equation of a line with 2 as X-intercept, 3 as Y-intercept.

Ans- a=2, b=3

$$\frac{x}{2} + \frac{y}{3} = 1$$

$$3x + 2y - 6 = 0$$

#### 5. Normal (Perpendicular) Form:-

The equation of a line L which is at a distance d from the origin.



$$x \cos \alpha + y \sin \alpha = d, d = |OP|$$

Ex:- find the equation of line which is at a distance 4 from origin and makes an angle  $135^\circ$  with respect to X-axis.

Here  $\alpha = 45^\circ; d = 4$

Therefore equation of line  $x \cos 45^\circ + y \sin 45^\circ = 4$

$$\Rightarrow x \frac{1}{\sqrt{2}} + y \frac{1}{\sqrt{2}} = 4 \Rightarrow \frac{1}{\sqrt{2}} (x + y) = 4 \Rightarrow x + y = 4\sqrt{2}$$



q-reduce  $x + \sqrt{3}y + 8 = 0$  to normal form of equation of straight line.  
 General form of eq of a line  $ax+by+c=0$

Normal form  $x \cos \alpha + y \sin \alpha - p = 0$

We know by comparing these two equation we get

$$\frac{a}{\sqrt{a^2 + b^2}}x + \frac{b}{\sqrt{a^2 + b^2}}y = \frac{-c}{\sqrt{a^2 + b^2}}$$

$$x + \sqrt{3}y + 8 = 0 \text{ here } a = 1, b = \sqrt{3}, c = 8$$

So equation in normal form will be

$$\frac{1}{\sqrt{1^2 + \sqrt{3}^2}}x + \frac{\sqrt{3}}{\sqrt{1^2 + \sqrt{3}^2}}y = \frac{-8}{\sqrt{1^2 + \sqrt{3}^2}}$$

$$\frac{1}{2}x + \frac{\sqrt{3}}{2}y = -\frac{8}{2}$$

$$\cos \frac{\pi}{3}x + \sin \frac{\pi}{6}y = -4$$

ex-find the equation of the line with normal of length 4 unit and inclined with an angle  $60^\circ$

$$\alpha = 60^\circ, p = 4$$

$$x \cos \alpha + y \sin \alpha = p$$

$$x \cos 60^\circ + y \sin 60^\circ = 4$$

$$x\left(\frac{1}{2}\right) + y\left(\frac{\sqrt{3}}{2}\right) = 4$$

$$x + \sqrt{3}y = 8$$

**General form of a Equation of a line**

$$**ax+by+c=0**$$

$$by+c = -ax$$

$$by = -ax - c$$

$$y = \frac{-a}{b}x - \frac{c}{b}, (\text{slope})m = \frac{-a}{b} \quad C(\text{y intercept}) = -\frac{c}{b}$$

ex- Find slope and y-intercept of the equation  $2x+3y+8=$

$$2x+3y+8=0$$

$$3y = -2x - 8$$

$$y = \frac{-2}{3}x + \left(-\frac{8}{3}\right)$$

$$m = \frac{-2}{3} \quad c = \left(-\frac{8}{3}\right)$$

Ex-Find angle of inclination of the equation  $2x+3y+8=0$

$$2x+3y+8=0$$

$$3y = -2x-8$$

$$y = \frac{-2}{3}x + \left(-\frac{8}{3}\right)$$

$$m = \frac{-2}{3} \quad \tan\theta = \frac{-2}{3}$$

$$\theta = \tan^{-1}\left(\frac{-2}{3}\right)$$

Ex-Find angle of inclination of the equation  $2x-2y+8=0$

$$2x-2y+8=0$$

$$-2y = -2x-8$$

$$y = \frac{2}{2}x + \left(\frac{8}{2}\right)$$

$$y = x+4$$

$$m=1, \tan\theta=1$$

$$\theta = \tan^{-1} 1 = 45^\circ$$

1. **find the equation** of a line which passes through (1,-1) and having **inclination  $150^\circ$**

What u think:-

What to do	What is given	Thinking's
Find Equation	Passes through (1,-1)	1. Slope-intercept form



of a line		2. Slope-Point form 3. Two point form 4. Intercept form 5. Normal form
	Angle of inclination $150^\circ$	Slope = $\tan \theta$ Slope = $\tan 150^\circ = -\frac{1}{\sqrt{3}}$

Find the equation of line passing through (1,-2) and having Y-intercept as 2.

What to do	What is given	Thinking's
Find Equation of a line	Passes through (1,-2)	1. Slope-intercept form 2. Slope-Point form 3. Two point form 4. Intercept form 5. Normal form
	Y-intercept:-2	As it has 2 as Y-intercept. Hence the line passes through point (0,2).

### PROBLEM ON THESE CONCEPT DISCUSSED EARLIER

$$\theta = \tan^{-1}\left(\frac{m_1 - m_2}{1 + m_1 m_2}\right)$$

$m_1 = m_2$  parallel

$m_1 m_2 = -1$  perpendicular

EX- Find angle between two lines  $x+y+6=0$  and  $2x-y+3=0$

Slope of the line  $x+y+6=0$  is

$$y = -x - 6,$$

$$m_1 = -1$$

slope of the line  $2x-y+3=0$

$$y = 2x + 3$$

$$m_2 = 2$$

$$\theta = \tan^{-1}\left(\frac{m_1 - m_2}{1 + m_1 m_2}\right)$$

$$\theta = \tan^{-1}\left(\frac{-1 - 2}{1 + (-1) \cdot 2}\right)$$

$$\theta = \tan^{-1}\left(\frac{-3}{-1}\right)$$

$$\theta = \tan^{-1}(3)$$

**ex- find the line perpendicular to  $x+y+2=0$  whose y-intercept is 3**

let the equation of the line perpendicular to  $x+y+2=0$  is

$$y=mx+c$$

$$\text{given } c=3$$

$$y=mx+3$$

since two lines are perpendicular then product of their slope is -1

slope of  $x+y+2=0$  is -1

$$(y = -x - 2)$$

$$m \cdot (-1) = -1$$

$$m = -1 / -1 = 1$$

so equation of the line is  $y=x+3$

$$x-y+3=0 \text{ (ans)}$$

**ex- find the line parallel to  $x+y+2=0$  whose y-intercept is 3**

let the equation of the line parallel to  $x+y+2=0$  is

$$y=mx+c$$

$$\text{given } c=3$$

$$y=mx+3$$

if two lines are parallel the slope are equal to each other

Then slope of the line  $x+y+2=0$  ( $y = -x - 2$ ) is -1

so equation of the line is  $y=-x+3$

$$x+y-3=0 \text{ (ans)}$$

**ex- check whether these two lines are perpendicular or parallel to each other.**

$$x+2y+3=0, 2x+4y+1=0$$

slope of the line  $x+2y+3=0$  is  $m_1 = \frac{-1}{2}$

$$2y = -x - 3$$

$$Y = -\frac{1}{2}x - \frac{3}{2}$$

slope of the line  $2x+4y+1=0$  is  $m_2 = \frac{-1}{2}$

$$4y = -2x - 1$$

$$y = -\frac{2}{4}x - \frac{1}{4}$$

$$m_1 = m_2$$

so these two lines are parallel

ex- check whether these two lines are perpendicular or parallel to each other.

$$x+2y+3=0, 4x-2y+1=0$$

slope of the line  $x+2y+3=0$  is  $m_1 = \frac{-1}{2}$

$$\theta = \tan^{-1}\left(\frac{m_1 - m_2}{1 + m_1 m_2}\right)$$

$$2y = -x - 3$$

$$\theta = \tan^{-1}\left(\frac{\frac{-1}{2} - 2}{1 + \frac{-1}{2} \cdot 2}\right)$$

$$Y = \frac{-1}{2}x - \frac{3}{2}$$

$$\theta = \tan^{-1}\left(\frac{-5}{0}\right) = \tan^{-1}(\infty) = 90^\circ$$

90°

slope of the line  $4x-2y+1=0$  is  $m_2 = 2$

$$2y = 4x + 1$$

$$y = \frac{4}{2}x + \frac{1}{2}$$

$$m_1 \cdot m_2 = \frac{-1}{2} \cdot 2 = -1$$

so these two lines are perpendicular

### Condition for coincidence of lines

$$.a_1x + b_1y + c_1 = 0$$

$$.a_2x + b_2y + c_2 = 0$$

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$$

Ex  $2x+3y+6=0$

$4x+6y+12=0$

$$\frac{2}{4} = \frac{3}{6} = \frac{6}{12} = \frac{1}{2}$$

### Equation of a line passing through a point and parallel or perpendicular to a line

Find the equation a line L


which passes through

\_\_\_\_\_  $(x_1, y_1)$  and parallel to \_\_\_\_\_ L

The equation of  $L_1$  is  $y = mx + c$

Slope of  $L_1 = m$

As  $L \parallel L_1$



Therefore slope of L= m

As L passes through  $(x_1, y_1)$  and having slope m therefore equation of line L in slope point form is  $y-y_1 = m(x-x_1)$

**Ex- find equation of a line which passes through a point(4,3) and parallel to the line  $2x-3y+5=0$**

Ans- passing through a poin  $(x_1, y_1)=(4,3)$

Line is parallel to  $2x-3y+5=0$  so slope will be equal to the slope of the given line.

Slope of  $2x-3y+5=0$

$$3y=2x+5$$

$$Y=\frac{2}{3}x + \frac{5}{3}$$

$$m = \frac{2}{3}$$

$$(2x-3y+k=0)$$

$$2.4-3.3+k=0$$

$$8-9+k=0$$

$$k=1, 2x-3y+1=0)$$

so equation of the line passing through  $(4,3)$  and slope  $\frac{2}{3}$

$$y-y_1 = m(x-x_1)$$

$$\text{is } y-3 = \frac{2}{3}(x-4)$$

$$3(y-3)=2(x-4)$$

$$3y-9=2x-8$$

$$2x-3y+1=0$$

**Equation of line passing through a point and perpendicular to a line:-**

Suppose L is a line

Which passes through  $(x_1, y_1)$  and perpendicular

To the line  $L_1: y=mx+c$


We know that if two Lines are perpendicular

Then the multiplication of Their slopes is -1.

Therefore slope of L X slope of  $L_1$

is -1.

Slope of  $L = -1/m$



Equation of line passing through  
( $x_1, y_1$ ) and having  $-1/m$  as slope is  
$$y - y_1 = -\frac{1}{m}(x - x_1)$$

**Ex2- find equation of a line which passes through a point(4,3) and perpendicular to the line  $2x-3y+5=0$**

Ans- passing through a poin ( $x_1, y_1$ )=(4,3)

Line is perpendicular to  $2x-3y+5=0$  so slope will be equal to  $-1/\text{slope of the given line}$ .

Slope of  $2x-3y+5=0$

$$3y=2x+5$$

$$Y=\frac{2}{3}x + \frac{5}{3}$$

$m = \frac{2}{3}$  slope of the line perpendicular to it=  $-\frac{3}{2}$

so equation of the line passing through (4,3) and slope  $-\frac{3}{2}$

$$\text{is } y-3 = \frac{-3}{2}(x-4)$$

$$2(y-3) = -3(x-4)$$

$$2y-6 = -3x+12$$

$$3x+2y-18=0(\text{ans})\text{eq of the line}$$

**Equation of a line passing through intersection of two lines**

If a line passing through intersection of two lines  $L_1$  and  $L_2$  (GIVEN)

Then the equation of the line is  $L_1 + kL_2=0$

**Obtain equation of the line passing through the intersection of  
 $3x-2y+7=0, x+3y+3=0$  and point (1, -1)**

Ans- equation of the line  $3x-2y+7+k(x+3y+3)=0$

$$3 \cdot 1 - 2(-1) + 7 + k(1 + 3(-1) + 3) = 0$$

$$3 + 2 + 7 + k(1 - 3 + 3) = 0$$

$$12 + k = 0$$

$$k = -12$$

So equation of the line is  $3x - 2y + 7 + k(x + 3y + 3) = 0$


$$3x - 2y + 7 - 12(x + 3y + 3) = 0$$

$$3x - 2y + 7 - 12x - 36y - 36 = 0$$

$$-9x - 38y + 29 = 0$$

$$9x + 38y - 29 = 0 \text{ (ans)}$$

### Perpendicular Distance of the point from a line



Let  $d$  be the distance from the point  $(x_1, y_1)$  to the line  $ax + by + c = 0$

$$d = \left| \frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}} \right|$$

ex- Find the length of the perpendicular drawn from the point  $(-5, 3)$  to the line  $3x + 4y - 6 = 0$

$$x_1 = -5, y_1 = 3$$

$$a = 3, b = 4, c = -6$$

$$d = \left| \frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}} \right|$$

$$d = \left| \frac{3(-5) + 4(3) - 6}{\sqrt{3^2 + 4^2}} \right|$$

$$d = \left| \frac{-15 + 12 - 6}{\sqrt{25}} \right| = \left| \frac{-9}{5} \right| = \frac{9}{5} \text{ (ans)}$$

If the area of triangle ABC is zero then the points A  $(x_1, y_1)$ ; B  $(x_2, y_2)$  & C  $(x_3, y_3)$  are collinear.

### Condition for coincidence of lines

$$.a_1x+b_1y+c_1=0$$

$$.a_2x+b_2y+c_2=0$$

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$$

$$\text{Ex } 2x+3y+6=0$$

$$4x+6y+12=0$$

$$\frac{2}{4} = \frac{3}{6} = \frac{6}{12} = \frac{1}{2}$$

## CIRCLE

**Circle:-** A circle is the locus of points which are equidistance from a fixed point. Here the fixed point is the centre of the circle and The fixed distance is called as the radius of the circle.

**Chord of a circle:-** The line segment joining any two pointson the circumference of the circle is called as the chord.

Here in the diagram CD is the chord.

**Diameter:-**The chord which passes through the centre of the circle is called as thediameter. Here EF is the diameter.

**Equation of the circle:-**

The equation the circle, whose centre is at (h,k) and the radius is r, is  $(x - h)^2 + (y - k)^2 = r^2$

If the centre is the origin i.e (0,0) then the equation of theCircle is  $x^2 + y^2 = r^2$

**Ex:- find the equation of the circle whose centre is at (2,3)and radius is 4.**

Solution:-

Here the centre of the circle is at (2,3)

Radius r = 4



Therefore the equation of circle is  $(x - 2)^2 + (y - 3)^2 = 4^2$

$$\Rightarrow x^2 + 4 - 2x + y^2 + 9 - 6y$$

$$= 16$$

$$\Rightarrow x^2 + y^2 - 2x - 6y - 3 = 0$$

**Ex:- find the equation of the circle whose centre is at (-1,4) and**



radius is 3 Solution:- Here the centre of the circle is at (-1,4)

Radius  $r = 3$

Therefore the equation of circle is  $(x - (-1))^2 + (y - 4)^2 = 3^2$

$$\Rightarrow (x + 1)^2 + (y - 4)^2 = 3^2$$

$$\Rightarrow x^2 + 1 + 2x + y^2 + 16 - 8y = 9$$

$$\Rightarrow x^2 + y^2 + 2x - 8y - 8 = 0$$

Ex:- Ex:- find the equation of the circle whose centre is at origin and radius is 1 Solution:-

### Equation of a circle passing through three points

**Example-** Find Equation of a circle which passes through (0,0)(1,0)(0,1)?

Ans-

We know equation of a circle

$$X^2 + y^2 + 2gx + 2fy + c = 0$$

As circle passes through (0,0)

$$0^2 + 0^2 + 2g \cdot 0 + 2f \cdot 0 + c = 0$$

$$C = 0 \text{-----(1)}$$

As circle passes through (1,0)

$$1^2 + 0^2 + 2g \cdot 1 + 2f \cdot 0 + c = 0$$

$$1 + 2g + C = 0 \text{-----(2)}$$

As circle passes through (0,1)

$$0^2 + 1^2 + 2g \cdot 0 + 2f \cdot 1 + c = 0$$

$$1 + 2f + C = 0 \text{-----(3)}$$

Now we will solve these three equation

$$C = 0 \text{-----(1)}$$

$$1 + 2g + C = 0 \text{-----(2)}$$

$$1 + 2f + C = 0 \text{-----(3)}$$

Putting  $c = 0$  in eq- 2 we get

$$1 + 2g = 0$$

$$2g = -1$$

$$g = \frac{-1}{2}$$

Putting  $c = 0$  in eq- 3 we get

$$1 + 2f = 0$$

$$2f = -1$$

$$f = \frac{-1}{2}$$

we get  $g = \frac{-1}{2}, f = \frac{-1}{2}, c = 0$




so equation of the circle is

$$x^2+y^2+2gx+2fy+c=0$$

$$x^2+y^2+2 \cdot \frac{-1}{2}x+2 \cdot \frac{-1}{2}y+0=0$$

$$x^2+y^2-x-y=0(\text{ans})$$

### 3dimensional geometry



Distance between two points  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$

$$d=\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

ex- Find distance between two points  $(1,2,4)$  and  $(-5,6,7)$ ?

Ans- we know distance between two points is

$$d=\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

here  $(x_1, y_1, z_1) = (1, 2, 4)$   $(x_2, y_2, z_2) = (-5, 6, 7)$

$$d=\sqrt{(-5 - 1)^2 + (6 - 2)^2 + (7 - 4)^2}$$

$$d = \sqrt{36 + 16 + 9} = \sqrt{61}$$

ex- Find distance between two points (1,-2,4) and (-1,3,1)?

Ans- we know distance between two points is


$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

here  $(x_1, y_1, z_1) = (1, 2, 4)$ ,  $(x_2, y_2, z_2) = (-1, 3, 1)$

$$d = \sqrt{(-1 - 1)^2 + (3 - 2)^2 + (1 - 4)^2}$$

$$d = \sqrt{4 + 1 + 9} = \sqrt{14}$$

section formula



if a point  $(x, y, z)$  divide the line segment joining two points  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  in  $m:n$  ratio internally


$$\text{then the point } (x, y, z) = \left( \frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}, \frac{mz_2 + nz_1}{m+n} \right)$$

if a point  $(x, y, z)$  divide the line segment joining two points  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  in  $m:n$  ratio externally

$$\text{then the point } (x, y, z) = \left( \frac{mx_2 - nx_1}{m-n}, \frac{my_2 - ny_1}{m-n}, \frac{mz_2 - nz_1}{m-n} \right)$$

**Ex- Find the coordinates of the points that divide the line segment joining the points (2,3,-2) and (6,32) internally in the ratio 3:1?**

Ans-



A horizontal line segment is shown with a blue dot in the middle. The left endpoint is labeled (2, 3, -2) and the right endpoint is labeled (6, 3, 2). The ratio 3:1 is written above the dot.

$$(2, 3, -2) \quad 3:1 \quad (6, 3, 2)$$


We know if a point  $(x, y, z)$  divide the line sement joining two points  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2,)$  in  $m:n$  ratio internally

$$\text{then the point } (x, y, z) = \left( \frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}, \frac{mz_2 + nz_1}{m+n} \right)$$

$$\text{here } (x_1, y_1, z_1) = (2, 3, -2) \quad (x_2, y_2, z_2) = (6, 3, 2) \quad m=3 \quad n=1$$

$$(x, y, z) = \left( \frac{3 \cdot 6 + 1 \cdot 2}{3+1}, \frac{3 \cdot 3 + 1 \cdot 3}{3+1}, \frac{3 \cdot 2 + 1 \cdot (-2)}{3+1} \right) = \left( \frac{20}{4}, \frac{12}{4}, \frac{4}{4} \right) = (5, 3, 1)$$

Find the *ratio in which the point*  $(9, -11, 1)$  *divides the line segment joining the points*  $(1, 5, -3)$  *and*  $(3, 1, -2)$ .



A horizontal line segment is shown with a blue dot in the middle. The left endpoint is labeled (1, 5, -3) and the right endpoint is labeled (3, 1, -2). The ratio K:1 is written above the dot.

$$(1, 5, -3) \quad (9, -11, 1) K:1 \quad (3, 1, -2)$$

We know if a point  $(x, y, z)$  divide the line sement joining two points  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2,)$  in  $m:n$  ratio internally

$$\text{then the point } (x, y, z) = \left( \frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}, \frac{mz_2 + nz_1}{m+n} \right)$$

$$\text{here } (x_1, y_1, z_1) = (1, 5, -3) \quad (x_2, y_2, z_2) = (3, 1, -2) \quad m=k \quad n=1$$

$$(x, y, z) = \left( \frac{k \cdot 3 + 1 \cdot 1}{k+1}, \frac{k \cdot 1 + 1 \cdot 5}{k+1}, \frac{k \cdot (-2) + 1 \cdot (-3)}{k+1} \right) =$$

$$\left( \frac{3k + 1}{k + 1}, \frac{k + 5}{k + 1}, \frac{-2k - 3}{k + 1} \right) = (9, -11, 1)$$

$$\frac{3k + 1}{k + 1} = 9$$

$$3k + 1 = 9(k + 1)$$

$$3k + 1 = 9k + 9$$

$$6k + 8 = 0$$

$$k = -\frac{8}{6} = -\frac{4}{3}$$

Therefore the point divides the line in a ratio 4:3 externally.

Centroid of a triangle:

$$\left( \frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right)$$

Mid point formula

Midpoint(x,y,z) of the line segment joining two points  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  is


$$(x, y, z) = \left( \frac{x_2 + x_1}{2}, \frac{y_2 + y_1}{2}, \frac{z_2 + z_1}{2} \right)$$

*Question- What is the midpoint of the line joining the points  $(-3, 4, 7)$  and  $(9, 0, 3)$ ?*

*Ans-* Midpoint(x,y,z) of the line segment joining two points  $(x_1, y_1, z_1) = (-3, 4, 7)$  and  $(x_2, y_2, z_2) = (9, 0, 3)$  is

$$(x, y, z) = \left( \frac{9 + (-3)}{2}, \frac{0 + 4}{2}, \frac{3 + 7}{2} \right) = \left( \frac{6}{2}, \frac{4}{2}, \frac{10}{2} \right) = (3, 2, 5)$$

## DIRECTION COSINE AND DIRECTION RATIO



$$l = \cos \alpha$$

$$m = \cos \beta$$

$$n = \cos Y$$

(l,m,n) is direction cosine of OP.

Let a,b,c be direction ratio of a line and l,m,n be direction cosine then

$$\frac{l}{a} = \frac{m}{b} = \frac{n}{c} = k$$

$$l = ak, m = bk, n = ck$$

$$l^2 + m^2 + n^2 = 1 \text{ (proof not in syllabus)}$$

$$a^2 k^2 + b^2 k^2 + c^2 k^2 = 1$$

$$k^2 (a^2 + b^2 + c^2) = 1, k = \frac{1}{\sqrt{a^2 + b^2 + c^2}}$$

$$l = ak = \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$

$$m = bk = \frac{b}{\sqrt{a^2 + b^2 + c^2}}$$

$$n = ck = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

direction ratio of a line joining two points  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  is  $(x_2 - x_1, y_2 - y_1, z_2 - z_1)$

**EX- IF A LINE has direction ratio 2, -1, -2 . determine its direction cosine**

**Given a=2, b=-1, c=-2**

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{2}{\sqrt{2^2 + (-1)^2 + (-2)^2}} = \frac{2}{\sqrt{9}} = \frac{2}{3}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \frac{-1}{\sqrt{2^2 + (-1)^2 + (-2)^2}} = \frac{-1}{\sqrt{9}} = \frac{-1}{3}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \frac{-2}{\sqrt{2^2 + (-1)^2 + (-2)^2}} = \frac{-2}{\sqrt{9}} = \frac{-2}{3}$$

**ex- Find the direction cosines of the line passing through the two points (-2, 4, -5) and (1, 2, 3)?**

**ans= direction ratio of the line joining the two points (-2, 4, -5) and (1, 2, 3) is**  
 $1 - (-2), 2 - 4, 3 - (-5)$

$$= (3, -2, 8)$$

**This means (a,b,c)=(3,-2,8)**

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{3}{\sqrt{3^2 + (-2)^2 + (8)^2}} = \frac{3}{\sqrt{77}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \frac{-2}{\sqrt{3^2 + (-2)^2 + (8)^2}} = \frac{-2}{\sqrt{77}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \frac{8}{\sqrt{3^2 + (-2)^2 + (8)^2}} = \frac{8}{\sqrt{77}}$$

**ex- show that the points A(2,3,-4), B(1,-2,3) and C(3,8,-11) are collinear.**

**Ans Direction ratios of line joining A and B are**

**1-2,-2-3, 3-(-4) that is  $-1,-5,7=(a_1,b_1,c_1)$**

**Direction ratios of line joining B and C are**

**3-1,8+2,-11-3 that is  $2,10,-14=(a_2,b_2,c_2)$**

**two lines are collinear if the ration of their direction ration is same**

**that is  $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$**

$$\frac{-1}{2} = \frac{-5}{10} = \frac{7}{-14} = \frac{-1}{2}$$

**Angle between two lines**

**If two lines having direction ratio  $(a_1, b_1, c_1)$  and  $(a_2, b_2, c_2)$  given or direction cosine  $(l_1, m_1, n_1)$  and  $(l_2, m_2, n_2)$  given**

$$\cos\theta = l_1l_2 + m_1m_2 + n_1n_2$$

$$\cos\theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

**Condition of parallelism:**

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$$

$$\frac{l_1}{l_2} = \frac{m_1}{m_2} = \frac{n_1}{n_2}$$

**Condition of perpendicularity**

$$a_1 \cdot a_2 + b_1b_2 + c_1c_2 = 0$$

$$l_1 \cdot l_2 + m_1m_2 + n_1n_2 = 0$$

**Ex-find the acute angle between two lines whose direction ratios are 2,3,6 and 1,2,2 respectively.**

**Ans- here direction ratio is given so angle between two lines is**

$$\cos\theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

**Here  $a_1, b_1, c_1 = 2,3,6$   $a_2, b_2, c_2 = 1,2,2$**

$$\cos\theta = \frac{2 \cdot 1 + 3 \cdot 2 + 6 \cdot 2}{\sqrt{2^2 + 3^2 + 6^2} \sqrt{1^2 + 2^2 + 2^2}} = \frac{20}{\sqrt{49} \sqrt{9}} = \frac{20}{7 \cdot 3} = \frac{20}{21}$$

$$\theta = \cos^{-1} \left( \frac{20}{21} \right) \text{ (ans)}$$

**ex- Find angle between the lines whose direction cosines are**

$$\left( \frac{\sqrt{3}}{4}, \frac{1}{4}, \frac{\sqrt{3}}{2} \right) \left( \frac{\sqrt{3}}{4}, \frac{1}{4}, -\frac{\sqrt{3}}{2} \right)$$

**ans if we know the direction cosines then angle between two lines is**

$$\cos\theta = l_1 l_2 + m_1 m_2 + n_1 n_2$$

$$\text{Here } l_1, m_1, n_1 = \left( \frac{\sqrt{3}}{4}, \frac{1}{4}, \frac{\sqrt{3}}{2} \right) \quad l_2, m_2, n_2 = \left( \frac{\sqrt{3}}{4}, \frac{1}{4}, -\frac{\sqrt{3}}{2} \right)$$

$$\cos\theta = \frac{\sqrt{3}}{4} \cdot \frac{\sqrt{3}}{4} + \frac{1}{4} \cdot \frac{1}{4} + \frac{\sqrt{3}}{2} \cdot -\frac{\sqrt{3}}{2} = \frac{3}{16} + \frac{1}{16} - \frac{3}{4} = \frac{3+1-12}{16} = -\frac{8}{16} = -\frac{1}{2}$$

$$\cos\theta = -\frac{1}{2} = \cos 120^\circ$$

$$\theta = 120^\circ \text{ (ans)}$$

## PLANE

**Equation of a plane**

$$ax+by+cz+d=0$$

**equation of xy plane z=0**

**equation of yz plane x=0**

**equation of zx plane y=0**

equation of a plane through three non collinear points

suppose three non collinear points are  $(x_1, y_1, z_1), (x_2, y_2, z_2), (x_3, y_3, z_3)$

then equation of the plane

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix} = 0$$

Equation of plane in intercept form

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

ex- find the equation of the plane with intercepts 2,-1,5 on x,y,and z axis respectively

ans= we know equation of a plane having intercept a,b,c is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

$$\frac{x}{2} + \frac{y}{-1} + \frac{z}{5} = 1$$

$$5x - 10y + 2z - 10 = 0$$

Transformation of general equation of a plane to normal form

$$ax+by+cz+d=0$$

$$\text{normal form } lx+my+nz-p=0$$

$$\frac{l}{a} = \frac{m}{b} = \frac{n}{c} = -\frac{p}{d}$$

$$\frac{a}{\sqrt{a^2+b^2+c^2}}x + \frac{b}{\sqrt{a^2+b^2+c^2}}y + \frac{c}{\sqrt{a^2+b^2+c^2}}z = p$$

$$P = \frac{d}{\sqrt{a^2+b^2+c^2}}$$

ex- find direction cosine of the normal to the plane  $x-y+1=0$

direction ratio (a,b,c) is (1,-1,0)

$$\text{direction cosines are } l = \frac{a}{\sqrt{a^2+b^2+c^2}} = \frac{1}{\sqrt{1^2+(-1)^2+0^2}} = \frac{1}{\sqrt{2}}$$

$$m = \frac{b}{\sqrt{a^2+b^2+c^2}} = \frac{-1}{\sqrt{1^2+(-1)^2+0^2}} = \frac{-1}{\sqrt{2}}$$

$$n = \frac{c}{\sqrt{a^2+b^2+c^2}} = \frac{0}{\sqrt{1^2+(-1)^2+0^2}} = 0$$



## Angle between two planes

$$\cos\theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

Ex- find angle between the two planes  $2x+2y-3z=5$  and  $3x-3y+5z=3$

Ans= here  $(a_1, b_1, c_1) = (2, 2, -3)$   $(a_2, b_2, c_2) = (3, -3, 5)$

$$\text{We know } \cos\theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} = \frac{2 \cdot 3 + 2 \cdot (-3) + (-3) \cdot 5}{\sqrt{2^2 + 2^2 + (-3)^2} \sqrt{3^2 + (-3)^2 + 5^2}}$$

$$= \frac{6 - 6 - 15}{\sqrt{17}\sqrt{43}} = -\frac{15}{\sqrt{731}}$$

$$\theta = \cos^{-1} \left( -\frac{15}{\sqrt{731}} \right)$$

## Distance of a point from a plane

Distance from a point  $(x_1, y_1, z_1)$  to a line  $ax + by + cz + d = 0$

$$d = \left| \frac{ax_1 + by_1 + cz_1 + d}{\sqrt{a^2 + b^2 + c^2}} \right|$$

Ex- find the distance from the point  $(2, -3, -1)$  to the plane  $2x - 3y + 6z + 7 = 0$ .

Ans- we know Distance from a point  $(x_1, y_1, z_1)$  to a line  $ax + by + cz + d = 0$

$$d = \left| \frac{ax_1 + by_1 + cz_1 + d}{\sqrt{a^2 + b^2 + c^2}} \right|$$

here  $(x_1, y_1, z_1) = (2, -3, -1)$   $a=2, b=-3, c=6$

$$d = \left| \frac{2 \cdot 2 + (-3)(-3) + 6(-1) + 7}{\sqrt{2^2 + (-3)^2 + 6^2}} \right| = \left| \frac{4 + 9 - 6 + 7}{\sqrt{4 + 9 + 36}} \right| = \frac{14}{7} = 2$$

## Equation of a plane parallel to another plane and passing through a point

If equation of a plane  $ax + by + cz + d = 0$  then equation of plane parallel to this and pass through the point  $(x_1, y_1, z_1)$  is

$$ax + by + cz + k = 0 \text{-----eq-1}$$

$$ax_1 + by_1 + cz_1 + k = 0$$

K=? and put it in eq 1

Ex- Find the equation of the plane which passes through the point (1,-1,4) and is parallel to the plane  $2x-3y+7z-11=0$

Ans : the equation of plane parallel to  $2x-3y+7z-11=0$  is

$$2x-3y+7z+k=0\text{-----}(1)$$

But it is given that the plane passes through (1,-1,4)

$$\text{So } 2.1-3(-1)+7.4+k=0$$

$$2+3+28+k=0$$

$$K=-33$$

Now we will put  $k=-33$  in eq-----(1)

$$2x-3y+7z-33=0$$

Therefore equation of the plane parallel to  $2x-3y+7z-11=0$  and passes through(1,-1,4) is  $2x-3y+7z-33=0$

Equation of a plane perpendicular to another plane and passing through a point

If equation of a plane  $ax+by+cz+d=0$  then equation of plane perpendicular to this and pass through the point  $(x_1, y_1, z_1)$  is

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

Ex- Find the equation of the plane which passes through (4,-2,1) and is perpendicular to the line whose direction ratio are 7,2,-3

Ans-

Equation of the plane passes through (4,-2,1) is

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

$$a(x - 4) + b(y - (-2)) + c(z - 1) = 0$$

Here  $a=7, b=2, c=-3$

$$7(x - 4) + 2(y - (-2)) + (-3)(z - 1) = 0$$

$$7x - 28 + 2y + 4 - 3z + 3 = 0$$

$$7x + 2y - 3z - 21 = 0$$

Therefore  $7x + 2y - 3z - 21 = 0$  is the required equation of the plane.

Plane through the intersection of two given plane

If equation of the given plane are

$$P_1: a_1x + b_1y + c_1z + d_1 = 0$$

$$P_2: a_2x + b_2y + c_2z + d_2 = 0$$

Then the new plane will be

$$P_1 + kP_2 = 0, k \text{ is a constant.}$$

**Ex- Find the equation of the plane which is perpendicular to the plane  $5x+3y+6z+8=0$  and contains the line of intersection of the planes  $x+2y+3z-4=0$  &  $2x+y-z+5=0$ .**

Ans- equation of the plane through the line of the intersection of the planes  $x+2y+3z-4=0$  &  $2x+y-z+5=0$ . is

$$(x+2y+3z-4)+k(2x+y-z+5)=0$$

$$(1+2k)x+(2+k)y+(3-k)z+(5k-4)=0 \text{-----eq-1}$$

Since the plane is perpendicular to  $5x+3y+6z+8=0$

So  $(1+2k)5+(2+k)3+(3-k)6=0$  (by using perpendicular condition)

$$5+10k+6+3k+18-6k=0$$

$$7k+29=0$$

$$K = \frac{-29}{7}$$

By putting  $k = \frac{-29}{7}$  in eq----1 we get

$$(1+2 \cdot \frac{-29}{7})x+(2+(\frac{-29}{7}))y+(3-(\frac{-29}{7}))z+(5(\frac{-29}{7})-4)=0$$

$$\left(1 - \frac{58}{7}\right)x + \left(2 - \frac{29}{7}\right)y + \left(3 + \frac{29}{7}\right)z + 5\left(-\frac{29}{7}\right) - 4 = 0$$

$$51x + 15y - 50z + 173 = 0$$

Therefore  $51x + 15y - 50z + 173 = 0$  is required equation of the plane.

## SPHERE

THE LOCOUS OF ALL THE POINT IN A SPACE WHICH ARE EQUIDISTANCE FROM A FIXED POINT IS KNOWN AS A SPHERE.

The fixed point is known as centre of the sphere and the constant distance is known as radius of the sphere.

Equation of a sphere whose centre at origin

$$(x - 0)^2 + (y - 0)^2 + (z - 0)^2 = r^2$$
$$x^2 + y^2 + z^2 = r^2$$

General equation of a sphere

Equation of a sphere having centre at  $(a, b, c)$  is  $(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2$

$$x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$$

Where center of the sphere is  $(-u, -v, -w)$  and radius  $r = \sqrt{u^2 + v^2 + w^2 - d}$

Equation of a sphere having co-ordinates of end point of diameter  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$

$$(x - x_1)(x - x_2) + (y - y_1)(y - y_2) + (z - z_1)(z - z_2) = 0$$

Ex- Find the centre and radius of a sphere

$$4x^2 + 4y^2 + 4z^2 - 16x - 24z + 3 = 0$$

Ans= We know general equation of a sphere is

$$x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0 \text{-----eq-1}$$

Given equation of a sphere is

$$4x^2 + 4y^2 + 4z^2 - 16x - 24z + 3 = 0$$

$$x^2 + y^2 + z^2 - 4x - 6z + \frac{3}{4} = 0 \text{-----eq 2}$$

$$x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0 \text{-----eq-1}$$

By comparing eq-1 and eq-2

$$2u = -4 \quad 2v = 0, \quad 2w = -6 \quad d = \frac{3}{4}$$

$$u = -2, \quad v = 0, \quad w = -3, \quad d = \frac{3}{4}$$

Centre of the sphere is  $(-u, -v, -w) = (2, 0, 3)$

Radius of the sphere is  $r = \sqrt{u^2 + v^2 + w^2 - d}$

$$r = \sqrt{u^2 + v^2 + w^2 - d}$$

$$r = \sqrt{(-2)^2 + 0^2 + (-3)^2 - \frac{3}{4}} = \sqrt{4 + 9 - \frac{3}{4}} = \sqrt{\frac{49}{4}} = \frac{7}{2}$$

Hence the centre is  $(2, 0, 3)$  and radius is  $7/2$ .

ex- Find the centre and radius of a sphere

$$x^2 + y^2 + z^2 - 2x - 4y - 6z - 11 = 0$$

We know general equation of a sphere is

$$x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0 \text{-----eq-1}$$

Given equation of a sphere is

$$x^2 + y^2 + z^2 - 2x - 4y - 6z - 11 = 0 \text{-----eq-2}$$

$$x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0 \text{-----eq-1}$$

By comparing eq-1 and eq-2

$$2u = -2 \quad 2v = -4, \quad 2w = -6 \quad d = -11$$

$$u = -1, \quad v = -2, \quad w = -3, \quad d = -11$$

Centre of the sphere is  $(-u, -v, -w) = (1, 2, 3)$

Radius of the sphere is  $r = \sqrt{u^2 + v^2 + w^2 - d}$

$$r = \sqrt{u^2 + v^2 + w^2 - d}$$

$$r = \sqrt{(-1)^2 + (-2)^2 + (-3)^2 - (-11)} = \sqrt{1 + 4 + 9 + 11} = \sqrt{25} = 5$$

Hence the centre is (1,2,3) and radius is 5.

**Ex- Find equation of the sphere on the join of (2,3,5) and (4,9,-3) as diameter.**

Equation of a sphere having co-ordinates of end point of diameter  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$

$$(x - x_1)(x - x_2) + (y - y_1)(y - y_2) + (z - z_1)(z - z_2) = 0$$

$$(x_1, y_1, z_1) = (2, 3, 5) \quad (x_2, y_2, z_2) = (4, 9, -3)$$

**Equation of the sphere is**

$$(x - 2)(x - 4) + (y - 3)(y - 9) + (z - 5)(z - (-3)) = 0$$

$$x^2 + y^2 + z^2 - 6x - 12y - 2z + 20 = 0$$

hence  $x^2 + y^2 + z^2 - 6x - 12y - 2z + 20 = 0$  is the required equation of the sphere.

### EQUATION OF A SPHERE PASSING THROUGH FOUR POINTS

EX- Find equation of a sphere which passes through the points (0,0,0), (1,0,0), (0,1,0) and (0,0,1)

Ans-

Let equation of the sphere be

$$x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0 \text{-----1}$$

Since (0,0,0) lies on eq-1

$$\text{So } 0^2 + 0^2 + 0^2 + 2u \cdot 0 + 2v \cdot 0 + 2w \cdot 0 + d = 0$$

$$d = 0 \text{-----2}$$

Since (0,1,0) lies on eq-1

$$0^2 + 1^2 + 0^2 + 2u \cdot 0 + 2v \cdot 1 + 2w \cdot 0 + d = 0$$

$$1 + 2v + d = 0 \text{-----eq3}$$

Since (0,0,1) lies on eq-1

$$0^2 + 0^2 + 1^2 + 2u \cdot 0 + 2v \cdot 0 + 2w \cdot 1 + d = 0$$

$$1 + 2w + d = 0 \text{-----eq4}$$

Since (1,0,0) lies on eq-1

$$1^2 + 0^2 + 0^2 + 2u \cdot 1 + 2v \cdot 0 + 2w \cdot 0 + d = 0$$

$$1 + 2u + d = 0 \text{-----eq5}$$

$$d = 0 \text{-----2}$$

$$1 + 2v + d = 0 \text{-----eq3}$$

$$1 + 2w + d = 0 \text{-----eq4}$$

$$1 + 2u + d = 0 \text{-----eq5}$$

By solving eq-2 to eq-4 we will get

$$d = 0, u = -\frac{1}{2}, v = -\frac{1}{2}, w = -\frac{1}{2}$$

Now we get the equation of sphere by putting value of u,v,w and d in equation 1

$$x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$$

$$x^2 + y^2 + z^2 + 2\left(-\frac{1}{2}\right)x + 2\left(-\frac{1}{2}\right)y + 2\left(-\frac{1}{2}\right)z + 0 = 0$$

$$x^2 + y^2 + z^2 - x - y - z = 0$$

**EX--** Find equation of a sphere which passes through the points (1,0,0), (0,1,0) and (0,0,1) AND WHOSE CENTRE LIES ON THE PLANE  $3X - Y + Z = 2$

$$3(-u) - (-v) + (-w) = 0$$

$$-3u + v - w = 0$$

$$1 + 2w + d = 0$$

$$1 + 2u + d = 0$$

$$1 + 2v + d = 0$$

**Ex-** Find the Equation of the sphere whose centre is on the point (1,2,3) and which touches the plane  $3x + 2y + z + 4 = 0$

**Ans=**

**Since the sphere touches the plane  $3x + 2y + z + 4 = 0$**

**Its radius = length of perpendicular from its centre (1,2,3) to the plane  $3x + 2y + z + 4 = 0$**

$$d = \left| \frac{3 \cdot 1 + 2 \cdot 2 + 1 \cdot 3 + 4}{\sqrt{3^2 + 2^2 + 1^2}} \right| = \frac{14}{\sqrt{14}} = \sqrt{14}$$

the required equation of the sphere is  $(x - 1)^2 + (y - 2)^2 + (z - 3)^2 = \sqrt{14}^2$

$$(x - 1)^2 + (y - 2)^2 + (z - 3)^2 = 14$$

$$x^2 + y^2 + z^2 - 2x - 4y - 6z = 0$$

G.P. GAJAPATI